

PCTWORLD INTELLECTUAL PROPERTY ORGANIZATION
International Bureau

INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁶ : C07K 5/062, A61K 38/05	A1	(11) International Publication Number: WO 96/16080
		(43) International Publication Date: 30 May 1996 (30.05.96)

(21) International Application Number: **PCT/US95/14989**(22) International Filing Date: **17 November 1995 (17.11.95)**(30) Priority Data:
08/345,820 21 November 1994 (21.11.94) US(71) Applicant: **CORTECH, INC. [US/US]; 6850 North Broadway,
Denver, CO 80221 (US).**(72) Inventors: **GYORKOS, Albert; 11795 Decatur Drive, West-
minster, CO 80234 (US). SPRUCE, Lyle, W.; 7747 Hoyt
Court, Arvada, CO 80005 (US).**(74) Agent: **VIKSNINS, Ann, S.; Schwegman, Lundberg & Woess-
ner, P.O. Box 2938, Minneapolis, MN 55402 (US).**(81) Designated States: **AL, AM, AT, AU, BB, BG, BR, BY, CA,
CH, CN, CZ, DE, DK, EE, ES, FI, GB, GE, HU, IS, JP,
KE, KG, KP, KR, KZ, LK, LR, LS, LT, LU, LV, MD,
MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD,
SE, SG, SI, SK, TJ, TM, TT, UA, UG, UZ, VN, European
patent (AT, BE, CH, DE, DK, ES, FR, GB, GR, IE, IT, LU,
MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM,
GA, GN, ML, MR, NE, SN, TD, TG), ARIPO patent (KE,
LS, MW, SD, SZ, UG).****Published***With international search report.**Before the expiration of the time limit for amending the
claims and to be republished in the event of the receipt of
amendments.***Schwegman, Lundberg &
Woessner, P.A.****JUN 12 1996****RECEIVED**(54) Title: **HUMAN NEUTROPHIL ELASTASE INHIBITORS**

(57) Abstract

The present invention relates to certain substituted oxadiazole, thiadiazole and triazole peptoids which are useful as inhibitors of serine proteases including human neutrophil elastase, equivalently known as human leukocyte elastase.

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AT	Austria	GB	United Kingdom	MR	Mauritania
AU	Australia	GE	Georgia	MW	Malawi
BB	Barbados	GN	Guinea	NE	Niger
BE	Belgium	GR	Greece	NL	Netherlands
BF	Burkina Faso	HU	Hungary	NO	Norway
BG	Bulgaria	IE	Ireland	NZ	New Zealand
BJ	Benin	IT	Italy	PL	Poland
BR	Brazil	JP	Japan	PT	Portugal
BY	Belarus	KE	Kenya	RO	Romania
CA	Canada	KG	Kyrgyzstan	RU	Russian Federation
CF	Central African Republic	KP	Democratic People's Republic of Korea	SD	Sudan
CG	Congo	KR	Republic of Korea	SE	Sweden
CH	Switzerland	KZ	Kazakhstan	SI	Slovenia
CI	Côte d'Ivoire	LI	Liechtenstein	SK	Slovakia
CM	Cameroon	LK	Sri Lanka	SN	Senegal
CN	China	LU	Luxembourg	TD	Chad
CS	Czechoslovakia	LV	Latvia	TG	Togo
CZ	Czech Republic	MC	Monaco	TJ	Tajikistan
DE	Germany	MD	Republic of Moldova	TT	Trinidad and Tobago
DK	Denmark	MG	Madagascar	UA	Ukraine
ES	Spain	ML	Mali	US	United States of America
FI	Finland	MN	Mongolia	UZ	Uzbekistan
FR	France			VN	Viet Nam
GA	Gabon				

HUMAN NEUTROPHIL ELASTASE INHIBITORS

5 The present invention relates to certain substituted oxadiazole, thiadiazole and triazole peptoids which are useful as inhibitors of serine proteases including human neutrophil elastase, equivalently known as human leukocyte elastase.

10

Background of the Invention

Human neutrophil elastase (HNE) is a proteolytic enzyme secreted by polymorphonuclear leukocytes (PMNs) in response to a variety of inflammatory stimuli.

15 This release of HNE and its extracellular proteolytic activity are highly regulated and are normal, beneficial functions of PMNs. The degradative capacity of HNE, under normal circumstances, is modulated by relatively high plasma concentrations of

20 α_1 -proteinase inhibitor (α_1 -PI). However, stimulated PMNs produce a burst of active oxygen metabolites, some of which (hypochlorous acid for example) are capable of oxidizing a critical methionine residue in α_1 -PI. Oxidized α_1 -PI has been shown to have limited

25 potency as an HNE inhibitor and it has been proposed that alteration of this protease/antiprotease balance permits HNE to perform its degradative functions in localized and controlled environments.

Despite this balance of protease/antiprotease

30 activity, there are several human disease states in which a breakdown of this control mechanism is implicated in the pathogenesis of the condition. Improper modulation of HNE activity has been suggested as a contributing factor in adult respiratory distress

35 syndrome, septic shock and multiple organ failure. A series of studies also have indicated the involvement of PMNs and neutrophil elastase in myocardial ischemia-reperfusion injury. Humans with below-normal levels of α_1 -PI have an increased probability of


developing emphysema. HNE-mediated processes are implicated in other conditions such as arthritis, periodontal disease glomerulonephritis, dermatitis, psoriasis, cystic fibrosis, chronic bronchitis, atherosclerosis, alzheimers disease, organ transplantation, corneal ulcers, and invasion behavior of malignant tumors.

There is a need for effective inhibitors of HNE as therapeutic and as prophylactic agents for the treatment and/or prevention of elastase-mediated problems.

Summary of the Invention

The principal object of the present invention is to provide certain new compounds which are useful as serine protease inhibitors, including human neutrophil elastase. These compounds are characterized by their relatively low molecular weight, high potency and selectivity with respect to HNE. They can be used effectively to prevent, alleviate or otherwise treat disease states characterized by the degradation of connective tissue by proteases in humans.


The novel compounds of the invention may be structurally illustrated by the following Formula (A):



25


Where Z is a carbonyl containing group, preferably an amino-carbonyl-containing, group X and Y are independently O, S or N, provided that at least one of X and Y is N, and R₁ is alkyl, haloalkyl, alkenyl,

haloalkenyl, alkynyl, phenyl, phenylalkenyl,
phenyalkyl or a heteroaryl group. Typically, Z is,



or any other equivalent carbonyl-containing moiety
5 which does not effect the ability of the compound to
inhibit serine proteases; and R₂ is a substituted or
unsubstituted alkyl, alkoxy, alkythio, phenyl or
cycloalkyl and R₃ is a carbonyl-containing moiety; or
Z is any other equivalent carbonyl-containing moiety
10 which does not affect the ability of the compound to
inhibit serine proteases.

The invention also contemplates intermediates for
preparing the compounds of Formula (A), the
intermediates being represented by Formula (B) as
15 follows:



where T is a strong acid that will protonate the free
amine and X, Y, R₁ and R₂ are as defined above.

Brief Description of the Drawings

Figures 1-3 illustrate the synthesis of compounds according to the invention. In particular,

Figure 1 describes the synthesis of 1-[(N-hydroxy)carboximidamido]-1-acetoxy-2-benzyloxycarbonylamino-3-(S)-methyl-butane;

Figure II describes the synthesis of 1-[3-[5-substituted]-1,2,4-oxadiazolyl]-1-acetoxy-2-benzyloxycarbonylamino-3-methyl-butane; and


Figure III describes the synthesis of (benzyloxycarbonyl)-L-valyl-N-[1-3-[5-substituted]-1,2,4-oxadiazolyl]carbonyl]-2-(S)-methylpropyl]-L-prolinamide.

Figure 4 illustrates substrate and enzyme subsite binding.

Detailed Description


As noted, the invention is directed to substituted heterocyclic compounds which have demonstrated activity against human elastase (HLE), also referred to as human neutrophil elastase (HNE).

The novel compounds of the invention may be structurally illustrated by the following Formula (A):



where Z is a carbonyl containing group, preferably an amino-carbonyl-containing, X and Y are independently O, S or N, provided that at least one of X and Y is N, and R₁ is alkyl, haloalkyl, alkenyl, haloalkenyl,


alkynyl, phenyl, phenylalkenyl, phenylalkyl or a heteroaryl group. Typically, Z is,



- 5 or any other equivalent carbonyl-containing moiety which does not effect the ability of the compound to inhibit serine proteases; and R₂ is substituted or unsubstituted alkyl, alkoxy, alkythio, phenyl or cycloalkyl and R₃ is a carbonyl-containing moiety; or
- 10 Z is any other equivalent carbonyl-containing moiety which does not affect the ability of the compound to inhibit serine proteases.


The invention also contemplates intermediates for preparing the compounds of Formula (A), the

15 intermediates being represented by Formula (B) as follows:



where T is a strong acid that will protonate the free amine and X, Y, R₁ and R₂ are as defined above.

The compounds of the invention may be further described as pseudopeptides substituted with (1, 2, 4) or (1, 3, 4) thia- or oxa-diazoles or triazoles as illustrated below:



wherein R₃ is defined above and further


R₁ is alkyl, alkenyl, haloalkenyl, alkynyl being
 10 linear or branched, a phenyl, phenylalkenyl, or
 phenylalkyl optionally substituted with halogen,
 cyano, nitro, haloalkyl, amino, aminoalkyl,
 dialkylamino, alkyl, alkenyl, alkynyl, alkoxy,
 haloalkoxy, carboxyl, carboalkoxy, alkylcarboxamide,
 15 arylcarboxamide, alkylthio, or haloalkylthio groups
 being linear or branched, a heteroaryl,
 heteroarylalkyl or heteroarylalkenyl wherein the
 heteroaryl groups being monocyclic five or six
 membered containing one or two heteroatoms such as
 20 oxygen, sulfur, or nitrogen in any combination and
 being optionally substituted with halogen, cyano,
 nitro, haloalkyl, amino, aminoalkyl, dialkylamino,
 alkyl, alkoxy, alkenyl, alkynyl, haloalkoxy, carboxyl,
 carboalkoxy, alkylcarboxamide, arylcarboxamide,
 25 alkylthio, or haloalkylthio groups being linear or
 branched.

X and Y are independently selected from oxygen or sulfur and nitrogen optionally substituted with alkyl, alkenyl, alkynyl being linear or branched, a phenyl,

phenylalkyl, or phenylalkenyl optionally substituted with halogen, cyano, nitro, haloalkyl, amino, aminoalkyl, dialkylamino, alkyl, alkenyl, alkynyl, alkoxy, haloalkoxy, carboxyl, carboalkoxy, alkylcarboxamide, arylcarboxamide, alkylthio, or haloalkylthio groups being linear or branched, a heteroaryl, heteroarylalkyl or heteroarylalkenyl wherein the heteroaryl groups being monocyclic five or six membered containing one or two heteroatoms such as oxygen, sulfur, or nitrogen in any combination and being optionally substituted with halogen, cyano, nitro, haloalkyl, amino, aminoalkyl, dialkylamino, alkyl, alkoxy, alkenyl, alkynyl, haloalkoxy, carboxyl, carboalkoxy, alkylcarboxamide, arylcarboxamide, alkylthio, or haloalkylthio groups being linear or branched so as to maintain the five membered aromatic heterocycle, provided that at least one of X or Y is nitrogen. It is also preferred that where X or Y is a substituted nitrogen, the other is not a substituted nitrogen.

R_2 is a linear or branched alkyl, alkylthio, alkylthioalkyl, alkoxy, cycloalkyl, alkylcycloalkyl, phenyl or phenylalkyl optionally substituted with terminal guanidine, halogen, cyano, nitro, hydroxy, haloalkyl, alkylthio, guanidine, alkyl guanidine, dialkyl guanidine or amidine.

In a particular embodiment of the invention, R_3 is the following Formula I;



30

R_1 is an optionally substituted phenylalkenyl, phenylalkyl, heteroarylalkyl or heteroarylalkenyl; Y

and X are selected from optionally substituted N, S, and O, provided that X or Y is N; and R₂ is 2-propyl, benzyl, 3-guanidinyl or 4-amidinylbenzyl.

The preferred R₂ substituents include isopropyl, benzyl, 3-guanidinyl and 4-amidinylbenzyl.

According to an additional embodiment of the invention, R₁ is an optionally substituted phenylalkyl; Y is O; X is N; and R₂ is 2-propyl.

The R₁ substituent is preferably chosen from the following: trifluoromethyl, methyl, difluoromethyl, 2,6-difluorobenzyl, benzyl, 2-phenylethyl, 3-methoxybenzyl, 3-phenylpropyl, 3-trifluoromethylbenzyl, 2-methoxybenzyl, 2-trifluoromethylphenylchloromethyl, 3-thienylmethyl, styryl, 4-trifluoromethylstyryl, 4-methoxystyryl, 4-methoxybenzyl, or phenyl.

As used herein, the term "optionally substituted" means, when substituted, mono to fully substituted.

As used herein, the term "alkyl" means C₁-C₁₅, however, preferably C₁-C₇.

As used herein, the term "alkenyl" means C₁-C₁₅, however, preferably C₁-C₇.

As used herein, the term "alkynyl" means C₁-C₁₅, however, preferably C₁-C₇.


As used herein, the term "aryl" is synonymous with "phenyl" and unless otherwise stated is optionally substituted.

The oxadiazoles of the present invention have been found to be potent inhibitors of the serine protease human neutrophil elastase (HNE). These compounds can be regarded as reversible inhibitors that presumably form a transition state intermediate with the active site serine residue. These oxadiazoles can be characterized by their low molecular weights, high selectivity with respect to HNE and stability regarding physiological conditions. Therefore, these compounds can be implemented to prevent, alleviate and/or otherwise treat diseases

which are mediated by the degradative effects associated with the presence of HNE. Their usage is of particular importance as they relate to various human treatment *in vivo* but may also be used as a
 5 diagnostic tool *in vitro*.






As described above, R_3 may also be a moiety which contributes to the oral activity of the compound. Techniques for conferring oral activity onto a compound are well known in the art. By way of example
 10 only, R_3 may be chosen from Formulas II-VI, described below:

Formula II




where X_1 is N or CH; X_2 is a group of the formula:

15





or,




- where W is H, or alkyl straight chained or branched;
 5 X_3 is Ala, bAla, Leu, Ile, Val, Nva, bVal, Met, or Nle
 or an N-methyl derivative or a bond; and X_4 is an
 amino acid consisting of proline, isoleucine, leucine,
 cyclohexylalanine, cysteine optionally substituted at
 the sulfur with alkyl, alkenyl, being linear or
 10 branched or phenyl optionally substituted with
 halogen, cyano, nitro, haloalkyl, amino, aminoalkyl,
 dialkylamino, alkyl, alkoxy, haloalkoxy, carboxyl,
 carboalkoxy, alkylcarboxamide, arylcarboxamide,
 alkylthio, haloalkylthio groups being linear or
 15 branched, a phenylalanine, indoline-2-carboxylic acid,
 tetrahydroisoquinoline-2-carboxylic acid optionally
 substituted with alkyl, alkenyl, haloalkenyl, alkynyl
 being linear or branched, halogen, cyano, nitro,
 haloalkyl, amino, aminoalkyl, dialkylamino, alkoxy,
 20 haloalkoxy, carbonyl, carboalkoxy, alkylcarboxamide,
 arylcarboxamide, alkylthio, or haloalkylthio group
 being linear or branched, tryptophan, valine,
 norvaline, norleucine, octahydroindole-2-carboxylic
 acid, lysine optionally substituted with H,
 25 alkylcarboxy or arylcarboxy, glycine optionally
 substituted at the nitrogen with hydrogen, alkyl,
 cycloalkyl, phenyl, bicycloalkyl, heteroaryl, alkenyl,
 alkynyl, cycloalkyl alkyl, bicycloalkyl alkyl,
 alkoxyalkyl, alkylthioalkyl, alkylaminoalkyl, fused
 30 aryl-cycloalkyl alkyl, fused heteroarylcycloalkyl,


dialkylaminoalkyl, carboxyalkyl or alkoxycarbonyl alkyl.

R_3 may also be:

Formula III,




Formula IV



5


or Formula V



wherein Z_2 and Z_5 are optionally substituted alkyl, aryl, or arylalkyl; Z_3 is optionally substituted alkyl, cycloalkyl, alkenyl, aryl, arylalkyl, an

aliphatic heterocycle, or aromatic heterocycle; Z_4 and Z_6 are hydrogen or methyl; A is selected from the group consisting of


5



and n is 0, 1 or 2;


In addition, R_3 may be represented by the following:

Formula VI



- 10 where G is nitrogen or carbon; J_1 and J_2 are independently substituted with hydrogen, alkyl, cycloalkyl, aryl, heteroaryl, alkylaryl or alkyheterolaryl and J_3 is selected from hydrogen, acyl
 15 aryl- SO_2 -, alkylaryl- SO_2 -, heterocycle- SO_2 -, alkyl-NH- SO_2 -, aryl-NH- SO_2 -, alkylaryl-NH- SO_2 -, or heterocycle-NH- SO_2 -. When G is carbon then this carbon may be substituted with hydrogen, alkyl, cycloalkyl, aryl, heteroaryl, alkylaryl or alkylaryl.

Another important embodiment of the invention is the provision of novel intermediates of the Formula (B) as follows:



5 where T is a strong acid that will protonate the free amine and X, Y, R₁ and R₂ are as defined above.

Methods for synthesizing the above mentioned R₃ constituents are described in EPA 0 529 568 A1 (Formula II); U.S. Patent 5,055,450 (Formula III-V);
 10 EPA 0 528 633; Warner et al., *J. Med. Chem.*, 37:3090-3099 (1994); Damewood et al., *J. Med. Chem.*, 37:3303-3312 (1994); Bernstein et al., *J. Med. Chem.*, 37:3313-3326 (1994) (Formula VI).

The compounds of the present invention are not
 15 limited to use for inhibition of human elastase. Elastase is a member of the class of enzymes known as serine proteases. This class also includes, for example, the enzymes chymotrypsin, cathepsin G, trypsin and thrombin. These proteases have in common
 20 a catalytic triad consisting of Serine-195, Histidine-57 and Aspartic acid-102 (chymotrypsin numbering system). The precise hydrogen bond network that exists between these amino acid residues allows the Serine-195 hydroxyl to form a tetrahedral intermediate
 25 with the carbonyl of an amide substrate. The decomposition of this intermediate results in the

release of a free amine and the acylated enzyme. In a subsequent step, this newly formed ester is hydrolyzed to give the native enzyme and the carboxylic acid. It is this carboxyl component that helps characterize the specificity for the enzyme. In the example in which the carboxyl component is a peptide, the alpha-substituent of the amino acid is predominately responsible for the specificity toward the enzyme. Utilizing the well accepted subsite nomenclature by Schechter and Berger (*Biochem. Biophys. Res. Commun.* 27:157(1967) and *Biochem. Biophys. Res. Commun.* 32:898 (1968)), the amino acid residues in the substrate that undergo the cleavage are defined as $P_1 \dots P_n$ toward the N-terminus and $P_1' \dots P_n'$ toward the C-terminus. Therefore, the scissile bond is between the P_1 and the P_1' residue of the peptide subunits. A similar nomenclature is utilized for the amino acid residues of the enzyme that make up the binding pockets accommodating the subunits of the substrate. The difference is that the binding pocket for the enzyme is designated by $S_1 \dots S_n$ instead of $P_1 \dots P_n$ as for the substrate (Figure 4).




Figure 4

The characteristics for the P_1 residue defining serine proteinase specificity is well established (see

Table 1). The proteinases may be segregated into three subclasses: elastases, chymases and tryptases based on these differences in the P₁ residues. The elastases prefer small aliphatic moieties such as
 5 valine whereas the chymases and tryptases prefer large aromatic hydrophobic and positively charged residues respectively.

One additional proteinase that does not fall into one of these categories is prolyl endopeptidase. The
 10 P₁ residue defining the specificity is a proline. This enzyme has been implicated in the progression of memory loss in Alzheimer's patients. Inhibitors consisting of α -keto heterocycles have recently been shown to inhibit prolyl endopeptidase; Tsutsumi et.
 15 al., *J. Med. Chem.* 37: 3492:3502 (1994). By way of extension, α -keto heterocycles as defined by Formula I allow for an increased binding in P' region of the enzyme.

20 Table 1. P₁ characteristics for Proteinase Specificity

PROTEINASE	REPRESENTATIVE	P ₁
Elastases	Human Neutrophil Elastase	small aliphatic residues
Chymases	alpha-Chymotrypsin, Cathepsin G	aromatic or large hydrophobic residues
25 Tryptases	Thrombin, Tryptin, Urokinase, Plasma Killikrein, Plasminogen Activator, Plasmin	positively charged residues
Other	Prolyl, Endopeptidase	proline

Since the P₁ residue predominately defines the
 30 specificity of the substrate, the present invention relates to P₁-P_n' modifications, specifically, certain alpha-substituted keto-heterocycles composed of 1,3,4 oxadiazoles, 1,2,4-oxadiazoles, 1,3,4-thiadiazoles,

1,2,4-thiadiazoles, 1-substituted, and 4-substituted 1,2,4-triazoles. By altering the alpha-substituent and the substituent on the heterocycle, the specificity of these compounds can be directed toward
5 the desired proteinase (e.g., small aliphatic groups for elastase).

The efficacy of the compounds for the treatment of various diseases can be determined by scientific methods which are known in the art. The following are
10 noted as examples for HNE mediated conditions:

- for acute respiratory distress syndrome, the method according to Human neutrophil elastase (HNE) model (AARD 141:227-677 (1990), or the Endotoxin induced acute lung injury model in minipigs (AARD
15 142:782-788 (1990)) may be used;

- in ischemia/reperfusion, the method according to the canine model of reperfusion injury (*J.Clin.Invest.* 81:624-629 (1988)) may be used.

The invention also provides pharmaceutical
20 compositions containing a compound of the formula herein and the process of synthesis which include the intermediates for the manufacturing of the invention.

The compounds of the present invention may be preferably formulated into suitable pharmaceutical
25 preparations such as tablets, capsules or elixirs, for oral administration or in sterile solutions or suspensions for parenteral administration. They can be administered to patients (human and animals) in need of such treatment in a dosage range of 5 to 500
30 mg per patient generally given several times, thus giving a total daily dose of from 5 to 2000 mg per day. The dose will vary depending on severity of disease, weight of patient and other factors.

The compounds may be formulated into
35 pharmaceutical compositions by compounding 10 to 500 mg of a compound or mixture thereof or of a physiologically acceptable salt with a physiologically acceptable vehicle, carrier, excipient, binder,

preservative, stabilizer, flavor, etc., in a unit dosage form as called for by accepted pharmaceutical practice. The amount of active substance in these compositions or preparations is such that a suitable
5 dosage in the range indicated is obtained.

Illustrative of the adjuvants which may be incorporated in tablets, capsules and the like are the following: a binder such as gum tragacanth, acacia, corn starch or gelatin; an excipient such as micro-
10 crystalline cellulose; a disintegrating agent such as corn starch, pregelatinized starch, alginic acid and the like; a lubricant such as magnesium stearate; a sweetening agent such as sucrose, lactose or saccharin; a flavoring agent such as peppermint, oil
15 of wintergreen or cherry. When the dosage unit form is a capsule, it may contain in addition to materials of the above type, a liquid carrier such as fatty oil. Various other materials may be present as coatings or to otherwise modify the physical form of the dosage
20 unit. For instance, tablets may be coated with shellac, sugar or both. A syrup or elixir may contain the active compound, sucrose as a sweetening agent, methyl and propyl parabens as preservatives, a dye and a flavoring such as cherry or orange flavor.

25 Sterile compositions for injection can be formulated according to conventional pharmaceutical practice by dissolving or suspending the active substance in a vehicle such as water for injection, a naturally occurring vegetable oil like sesame oil,
30 coconut oil, peanut oil, cottonseed oil, etc., or a synthetic fatty vehicle like ethyl oleate or the like. Buffers, preservatives, antioxidants and the like can be incorporated as required. Pharmaceutical compositions containing the compounds of this
35 invention can also be administered topically. This can be accomplished by simply preparing a solution of the compound to be administered, preferably using a solvent known to promote transdermal absorption such

as ethanol or dimethyl sulfoxide (DMSO) with or without other excipients. Preferably topical administration will be accomplished using a patch either of the reservoir and porous membrane type or of a solid matrix variety.

Some suitable transdermal devices are described in U.S. Pat. Nos. 3,742,951, 3,742,951, 3,797,494, 3,996,934, and 4,031,894. These devices generally contain a backing member which defines one of its face surfaces, an active agent permeable adhesive layer defining the other face surface and at least one reservoir containing the active agent interposed between the face surfaces. Alternatively, the active agent may be contained in a plurality of microcapsules distributed throughout the permeable adhesive layer. In either case, the active agent permeable adhesive, is in contact with the skin or mucosa of the recipient. If the active agent is absorbed through the skin, a controlled and predetermined flow of the active agent is administered to the recipient. In the case of microcapsules, the encapsulating agent may also function as the membrane.

In another device for transdermally administering the compounds in accordance with the present invention, the pharmaceutically active compound is contained in a matrix from which it is delivered in the desired gradual, constant and controlled rate. The matrix is permeable to the release of the compound through diffusion or microporous flow. The release is rate controlling. Such a system, which requires no membrane is described in U.S. Pat. No. 3,921,636. At least two types of release are possible in these systems. Release by diffusion occurs when the matrix is non-porous. The pharmaceutically effective compound dissolves in and diffuses through the matrix itself. Release by microporous flow occurs when the pharmaceutically effective compound is transported through a liquid phase in the pores of the matrix.

While the invention has been described in connection with specific embodiments thereof, it will be understood that it is capable of further modifications and this application is intended to cover any variations, uses, or adaptations of the invention following, in general, the principles of the invention and including such departures from the present disclosure as come within known or customary practice within the art to which the invention pertains and as may be applied to the essential features hereinbefore set forth, and as follows in the scope of the appended claims.

EXAMPLES

The compounds of the present invention, salts thereof, and their intermediates can be prepared or manufactured as described herein or by various process known to be present in the chemical art. By way of an example, the final step in the process defined herein necessitates an oxidation of a 2° alcohol to the ketone in Formula I. As described here, this transformation from alcohol to ketone was utilized by way of the procedure of N-chlorosuccinimide (NCS) and dimethylsulfide followed by treatment with base. However, alternative methods are known to perform a similar conversion that include dimethylsulfoxide and oxalyl chloride followed by base and 1,1,1-tria¹acetoxy-1,-dihydro-1,2-benziodoxol-3(1H)-one (Dess-Martin's Periodinane), to mention two. Other methods may also be appropriate as described in Oxidation in Organic Chemistry by Milos Hudlicky, ACS Monograph 186 (1990).

By way of further example, the R₂ group is introduced by the appropriate selection of the chiral amino alcohol, either commercially available or prepared by methods available to one skilled in the art by way of reduction of the natural or unnatural amino acid. The amino alcohol is selectively protected allowing for the alcohol to be converted to

the corresponding aldehyde. By way of an alternative process, an amino protected aldehyde may be obtained by a reductive method of an amide made from the desired nitrogen protected amino acid and N,O-

5 dimethylhydroxylamine as described by Winreb (*Tetrahedron Lett.* 22:3815 (1981)). The aldehyde is subsequently allowed to react with acetone cyanohydrin to give the amino protected cyanohydrin (see Figure I). The cyanohydrin may also be prepared by a

10 standard method using sodium or potassium cyanide. Alternatively, trimethylsilylcyanide may be used to effect the same conversion from aldehyde to cyanohydrin after acidic workup of the O-trimethylsilylcyanohydrin intermediate. The next step

15 is the protection of the cyanohydrin's alcohol followed by the conversion of the nitrile to the amide oxime through the use of hydroxylamine. This intermediate allows for the incorporation the R_1 group into the heterocycle through the reaction of the amide

20 oxime amine with an activated R_1 carboxylic acid. The carboxylic acid may be activated by way of an acid chloride or fluoride, anhydride or an active ester with the requirements of the appropriate activation dictated by the R_1 group. The resulting N-acylated

25 amide oxime can be either isolated or allowed to proceed in the cyclization when subjected to elevated temperatures. If the N-acylated amide oxime is isolated it can be characterized and subsequently cyclized in a separate step to the 1,2,4-oxadiazole

30 (see Figure II).

In an anticipated analogy, the 1,2,4-triazole is envisioned by the reaction of an N-substituted hydrazine with the nitrile described herein. After acylation at either available nitrogen, the resulting

35 intermediate is allowed to form the 1,3,5-tri-substituted-1,2,4-triazole (Formula I, Y = substituted N and X = N) under conditions similar to that required for the 1,2,4-oxadiazole.

As described in Figure III, the alcohol and amine are deprotected by methods available to one skilled in the art and the resulting amine is then subsequently coupled to the desired R_3 carboxylic acid. The final
5 step, as mentioned herein, is the conversion of the 2° alcohol to the ketone. This oxidation can be performed by a number of methods with one such method being the use of the intermediate resulting from the reaction between NCS and dimethyl sulfide with the 2°
10 alcohol followed by base. This process affords the ketone and preserves the stereochemistry of the neighboring R_2 group.

The following examples are given to illustrate the invention and are not intended to be inclusive in
15 any manner:

i) temperatures are given in degrees Celsius (C); room temperature ranged from 17° to 26°;

ii) chromatography was performed as described by Still (*J.Org.Chem.* 43:2923 (1978)) using ICN silica
20 gel (60 Å), thin layer chromatography was performed using silica gel 60 F254 (25mm) precoated plates from EM Science;

iii) NMR chemical shift data is reported in parts per million (ppm) relative to tertamethylsilane (TMS, 0.00 ppm) for ^1H and deuterated chloroform (CDCl_3 , 77.00 ppm) for ^{13}C ; the abbreviations used for the
25 peak shape for the ^1H data are as follows: s-singlet; d-doublet; t-triplet; q-quartet; m-multiplet; and br-broad;

iv) the course of reactions was monitored by TLC
30 using UV and/or staining by aqueous KMnO_4 as detection methods, yields and reaction times are presented as illustrations but should not be considered optimal;

v) evaporation of solvents was performed using a
35 rotary evaporator under reduced pressure ranging between 5 and 35 mm Hg; the bath temperatures were between 17 and 50°C;

vi) chemical symbols have the standard meanings as familiar to someone skilled in the art, by way of example the following have been used: mmol (millimoles), mol (mole), mL (milliliters), L (liters), mg (milligram), g (grams), min (minutes), h (hours), TLC (thin layer chromatography), R_f (ratio between the distance traveled by the compound compared to that of the solvent as related to TLC), RP-HPLC (reverse phase-high pressure liquid chromatography), RT (room temperature).

EXAMPLE I - Methods of Synthesis

(Benzyloxycarbonyl)-L-Valyl-N-[1-[3-[5-(3-Trifluoromethylbenzyl)-1,2,4-oxadiazolyl]carbonyl]-2-(S)-Methylpropyl]-L-Prolinamide.

To a stirred mixture of 0.68 g (5.1 mmol) of N-chlorosuccinimide in 15 mL of dry toluene at 0°C was added 0.53 mL (7.3 mmol) of dimethylsulfide (DMS) under a nitrogen atmosphere. A white precipitate formed after the addition of DMS. After 30 minutes, the resulting suspension was cooled to -25°C using a carbon tetrachloride and dry ice bath. A solution of (benzyloxycarbonyl)-L-valyl-N-[1-[3-[5-(3-trifluoromethylbenzyl)-1,2,4-oxadiazolyl]hydroxymethyl]-2-(S)-methylpropyl]-L-prolinamide (0.84 g, 1.27 mmol) in 20 mL of dry toluene, was added dropwise. The resulting mixture was stirred for 3 h at -25°C and 0.75 mL (5.4 mmol) of triethylamine was added. After 15 minutes, the cold bath was removed and the reaction monitored by TLC; silica gel; ethyl acetate: hexanes (4:1). After 1 h, the mixture was diluted with ethyl acetate and washed with NaHCO₃ (saturated), brine and dried with anhydrous magnesium sulfate. The resulting residue was purified by column chromatography; silica gel, ethyl acetate:hexanes, (1:1 to 7:3) to give a semisolid 91.7% pure. The material was further

purified via RP-HPLC isocratic $\text{CH}_3\text{CN}:\text{H}_2\text{O}$ (3:2), to give the title compound as a white solid after lyophilization; TLC: $R_f = 0.69$, ethyl acetate: hexane (4:1).

5 ^1H NMR (CDCl_3): δ [0.90 (d, $J = 6.7$ Hz); 0.96 (d, $J = 6.7$ Hz); 1.01 (d, $J = 6.7$ Hz); 1.02 (d, $J = 6.7$ Hz); 12H]; 1.80 - 2.20 (m, 4 H); 2.25 - 2.50 (m, 2H), 3.54 - 3.70 (m, 1H), 3.70 - 3.83 (m, 1 H); 4.30 - 4.40 (m, 1H), 4.38 (s, 2H); 4.63 (dd, $J_1 = 2.7$ Hz, $J_2 = 7.1$ Hz, 1H); 5.11 (ABq, $J = 12.3$ Hz, 2H), 5.29 (dd, $J_1 = 5.0$ Hz, $J_2 = 7.3$ Hz, 1H), 5.60 d, $J = 9.1$ Hz, 1H); 7.30 - 7.40 (m, 5H), 7.45 - 7.64 (m, 6H).

10 ^{13}C NMR (CDCl_3) δ 17.24, 17.59, 19.46, 19.81, 25.12, 27.08, 30.26, 31.43, 32.69, 47.78, 57.49, 59.70, 61.53, 66.96, 123.73 (q, $J = 272$ Hz), 124.93 (q, $J = 3.7$ Hz), 125.85 (q, $J = 3.8$ Hz), 128.01, 128.13, 128.51, 129.63, 131.48 (q, $J = 32.6$ Hz), 132.44, 133.41, 136.27, 156.42, 164.87, 171.10, 172.29, 178.65, 189.89.

20 IR (Deposit) 3429.2, 3299.0, 2968.5, 1719.2, 1680.3, 1632.1 cm^{-1} .



25	%C	%H	%N	%F
Theory	60.27	5.82	10.65	8.67
Found	60.04	6.03	10.60	8.08

The intermediate (Benzyloxycarbonyl)-L-Valyl-N-[1-[3-[5-(3-Trifluoromethylbenzyl)-1,2,4-oxadiazolyl]hydroxymethyl]-2-(S)-Methylpropyl]-L-Prolinamide was prepared as follows:

a. N-Benzyloxycarbonyl-L-Valinol.

35 To a stirred solution of valinol (13.96 g, 0.14 mol) and triethylamine (18.87 mL, 0.14 mol) in 250 mL of dry methylene chloride cooled to 0°C under a nitrogen atmosphere was added 19.3 mL (0.14 mol) of

benzyl chloroformate dropwise over 20 min. After warming to RT overnight, the mixture was diluted with methylene chloride (100 mL) and washed with 1 N HCl (2X), brine and dried with anhydrous magnesium sulfate. The resulting mixture was filtered and evaporated to give 30.70 g (95.6 %) of N-benzyloxycarbonyl-valinol as a white solid: $R_f = 0.27$ (silica gel; 1:1 hexane:ethyl acetate). Used without further purification.

10

b. N-Benzyloxycarbonyl-L-Valinal.

To a stirred solution of N-benzyloxycarbonyl-L-valinol (8.80 g, 0.037 mol) and triethylamine (30.00 g, 41.3 mL, 0.30 mol) in 100 mL dimethylsulfoxide under a nitrogen atmosphere was added a solution of sulfur trioxide pyridine complex (20.62 g, 0.130 mol) in dimethylsulfoxide (100 mL) dropwise via an addition funnel. An ice water bath was used to maintain the temperature of the reaction mixture below room temperature during the addition. After 30 min the cooling bath was removed and the reaction was allowed to warm to room temperature over 2 h. The brown solution was acidified with cold 2 N HCl to approximately pH 2 and extracted with ether (3 x 150 mL). The combined organic layers were washed with brine, dried (anhydrous magnesium sulfate) and evaporated to give 7.40 g of N-benzyloxycarbonyl-L-valinal (84.8 %) as a yellow oil. The resulting material can be used without purification or further purified using column chromatography (silica gel, 9:1 hexanes:ethyl acetate).

^1H NMR (CDCl_3) δ 0.95 (d, $J = 7.0$ Hz, 3H), 1.04 (d, $J = 7.0$ Hz, 3H), 2.20-2.40 (m, 1H), 4.34 (dd, $J_1 = 4.1$, $J_2 = 7.4$ Hz, 1 H), 5.12 (s, 2H), 5.37 (brs, 1H), 7.34 (s, 5H), 9.65 (s, 1H).

c. 3-(S)-[(Benzyloxycarbonyl)amino]-2-Hydroxy-4-Methylpentanenitrile.

To a solution containing N-benzyloxycarbonyl-L-valinal (22.2 g, 0.094 mol) in methylene chloride (500 mL) and triethylamine (5.81 g, 8.0 mL, 0.057 mol) was added acetone cyanohydrin (24.23 g, 26.0 mL, 0.2847 mol). The reaction was allowed to stir at room temperature overnight and then concentrated under reduced pressure. The residue was taken up into ether and washed with brine (3X). The solvent was removed under reduced pressure and purified by column chromatography (silica gel; 1:2 acetone:hexane) to give 21.38 g (86.6%) of 3-(S)-[(benzyloxycarbonyl)amino]-2-hydroxy-4-methylpentanenitrile as a pale yellow oil; TLC R_f = 0.33 (silica gel; 1:2 acetone:hexane).

^1H NMR (CDCl_3) δ 0.92-1.05 (m, 6H), [1.80-2.0 (m), 2.10-2.30 (m), 1H], [3.45-3.55 (m), 3.70-3.80 (m), 1H], 4.65 (m, 1 H), [5.14 (s), 5.15 (s), 2H], 5.35 (d, J = 4.8 Hz, 1H), 7.36 (s, 5H).

d. 3-(S)-[(Benzyloxycarbonyl)amino]-2-Acetoxy-4-Methylpentanenitrile.

To a solution containing 3-[(benzyloxycarbonyl)amino]-2-hydroxy-4-methylpentanenitrile (32.0 g, 0.12 mol) and pyridine (59 mL) was added acetic anhydride (73.6 g, 68 mL, 0.72 mol) dropwise at RT. After 3 h the reaction was diluted with ethyl acetate and washed with water. The organic layer was separated, dried (anhydrous magnesium sulfate) and evaporated. The residue was purified by column chromatography (silica gel, 1:1 hexane:ethyl acetate) to give 33.08 g (94.0 %) of 3-(S)-[(benzyloxycarbonyl)amino]-2-acetoxy-4-methylpentanenitrile as a viscous yellow oil; TLC R_f = 0.70 (silica gel, 1:1 hexane:ethyl acetate).


^1H NMR (CDCl_3) δ 0.95-1.09 (m, 6H), 1.82-2.20 (m, 1H), [2.06 (s), 2.09 (s), 3H], 3.86-4.04 (m, 1H), [4.90 (d, J = 8.5 Hz), 4.93 (d, J = 9.7 Hz), 1H], [5.12 (s), 5.14 (s), 5.15 (s), 5.16 (s), 2H], [5.48

(d, J = 4.9 Hz), 5.58 (d, J = 4.0 Hz), 1H], 7.36 (s, 5H).

- e. 1-[(N-Hydroxy)carboximidamido]-1-Acetoxy-2-(S)-
5 Benzyloxycarbonyl amino-3-Methyl-Butane.

A solution containing 3-(S)-
[(benzyloxycarbonyl)amino]-2-acetoxy-4-
methylpentanenitrile (33.0 g, 0.11 mol), hydroxylamine
hydrochloride (9.90 g, 0.14 mol) and sodium acetate
10 (13.2 g, 0.16 mol) in a mixture of ethanol (330 mL)
and water (66 mL) was heated to reflux under a
nitrogen atmosphere for 3 h. The reaction mixture
was concentrated, diluted with ethyl acetate and
washed with water. The organic layer was dried
15 (anhydrous magnesium sulfate), evaporated and purified
via column chromatography to yield 22.5 g (58.2 %) of
1-[(N-hydroxy)carboximidamido]-1-acetoxy-2-(S)-
benzyloxycarbonyl amino-3-methyl-butane as a yellow
foam. An analytical sample was prepared by RP-HPLC
20 (C₁₈, 2:3 acetonitrile:water with 0.1 % TFA,
isocratic). The pure fractions were combined and
evaporated. The residue was taken up into ethyl
acetate, washed with dilute sodium bicarbonate, dried
(anhydrous magnesium sulfate) and evaporated. The
25 resulting residue was dissolved in dry ether, cooled
to -78°C under a nitrogen atmosphere and 1.1
equivalents of 4.0 N HCl in dioxane was added. The
resulting white solid was collected by filtration and
dried.

30



	%C	%H	%N
Theory	51.41	6.47	11.24
Found	51.30	6.40	11.17

35

- f. 3-Trifluoromethyl phenylacetyl Chloride.

To a mixture of 10.00 g (48.96 mmol) of 3-
trifluoromethyl phenylacetic acid dissolved in 30 mL

of dry methylene chloride was added 6.4 mL (9.33 g, 73.44 mmol) of oxalyl chloride over 2 minutes. Two drops of DMF was added (gas evolution) and the reaction was allowed to stir for 2 h at RT (gas evolution stopped). The reaction mixture was evaporated (40°C, aspirator vacuum) to give (3-trifluoromethyl)phenylacetyl chloride as a yellowish oil, 10.74 g (98.5%). The product was used without further purification.

10 IR (neat) 1796.8 cm^{-1} C = O (acid chloride).

g. 1-[(N-Hydroxy)carboximid-N-(3-Trifluoromethylphenylacetyl)-amido]-1-Acetoxy-2-(S)-Benzyloxycarbonylamino-3-Methyl-Butane.

15 To 1-[(N-hydroxy)carboximidamido]-1-acetoxy-2-(S)-benzyloxycarbonyl-amino-3-methyl-butane (6.74 g, 20 mmol) was added 40 mL of dry toluene and 25 mL of chloroform under a nitrogen atmosphere. The solution was cooled in a brine/ice bath and 4.2 mL (30 mmol) of triethylamine was added. After 5 minutes, 3-trifluoromethyl phenylacetyl chloride (4.67 g, 21.0 mmol), dissolved in 25 mL of chloroform, was added dropwise over 10 minutes. The reaction mixture was allowed to warm to room temperature overnight. The reaction was determined to be complete by TLC (silica gel; ethyl acetate: hexane, 1:1). Visualization was realized by aqueous KMnO_4 staining, the amide oxime develops at RT whereas the product requires heating. The mixture was evaporated under reduced pressure and the residue purified by column chromatography (silica gel; ethyl acetate:hexane, 50:50 to 70:30) to give 4.08 g (40%) of 1-[(N-hydroxy)carboximid-N-(3-trifluoromethylphenylacetyl)amido]-1-acetoxy-2-(S)-benzyloxycarbonyl-amino-3-methyl-butane as an off-white solid.

^1H NMR (CDCl_3) δ [0.90 (d, J = 6.8 Hz); 0.95 (d, J = 6.8 Hz); 0.98 (d, J = 6.8 Hz); 1.02 (d, J = 6.8 Hz); 6H]; 1.82 - 2.00 (m, 1H), [1.95 (s); 2.09 (s);

3H]; 3.80 - 3.98 (m, 1H); [3.80 (s), 3.85 (s), 2H],
4.80 - 5.20 (m, 5H), 5.41 (t, J = 6.5 Hz, 1H); 7.30 -
7.40 (m, 5H); 7.40 - 7.60 (m, 4H).

- 5 h. 1-[3-[5-(3-Trifluoromethylbenzyl)]-1,2,4-
oxadiazolyl]-1-Acetoxy-2-(S)-
Benzyloxycarbonylamino-3-Methyl-Butane.

To 1-[(N-hydroxy)carboximid-N-(3-
trifluoromethylphenylacetyl) amido]-1-acetoxy-2-(S)-
10 benzyloxycarbonylamino-3-methyl-butane (4.08 g, 7.8
mmol), was added dry toluene (100 mL) under a nitrogen
atmosphere and heated to reflux for 60 h. The mixture
was cooled to room temperature and evaporated to give
an oil. This resulting material was purified by
15 column chromatography (silica gel; ethyl acetate:
hexane, 50:50 to 60:40) to give 3.62 g (91.8%) of 1-
[3-[5-(3-trifluoromethylbenzyl)]-1,2,4-oxadiazolyl]-1-
acetoxy-2-(S)-benzyloxycarbonylamino-3-methyl-butane
as an oil.

20 ¹H NMR (CDCl₃) δ [0.93 (d, J = 6.7 Hz); 1.00 (d,
J = 6.7 Hz); 1.04 (d, J = 6.7 Hz); 6H]; [1.54 - 1.68
(m); 1.73 - 1.87 (m), 1H]; [2.05 (s); 2.14 (s); 3H];
4.00 - 4.20 (m, 1H), [4.24 (s), 4.31 (s); 2H]; [4.99
(s), 5.09 (s), 2H]; [5.03 (d, J = 10.5 Hz), 5.37 (d, J
25 = 10.5 Hz); 1H], [6.08 (d, J = 1.8 Hz), 6.10 (d, J =
4.4 Hz), 1H], 7.28- 7.62 (m, 9H).

- i. 1-[3-[5-(3-Trifluoromethylbenzyl)]-1,2,4-
oxadiazolyl]-2-(S)- Benzyloxycarbonylamino-3-Methyl-
30 Butan-1-ol.

To 1-[3-[5-(3-trifluoromethylbenzyl)]-1,2,4-
oxadiazolyl]-1-acetoxy-2-(S)-benzyloxycarbonylamino-3-
methyl-butane (3.62 g, 7.16 mmol), was added 70 mL of
methanol and stirred until homogeneous (10 minutes).
35 To this resulting solution was added potassium
carbonate (1.19 g, 8.59 mmol) dissolved in 20 mL of
water. The reaction was monitored by TLC (silica gel;
ethyl acetate: hexane, 7:3). After 45 minutes, the

reaction was diluted with ethyl acetate and washed with water (2x). The organics were dried with anhydrous magnesium sulfate, filtered and the solvent removed under reduced pressure. The residue was
5 purified by column chromatography (silica gel; ethyl acetate: hexane, 50:50 to 60:40) to give 2.45 g (74.1%) of 1-[3-[5-(3-trifluoromethylbenzyl)]-1,2,4-oxadiazolyl]-2-(S)-benzyloxycarbonylamino-3-methylbutan-1-ol.

10 ¹H NMR (CDCl₃) δ [0.92 (d, J = 6.7 Hz); 0.96 (d, J = 6.7 Hz); 0.98 (d, J = 6.7 Hz); 1.06 (d, J = 6.7 Hz); 6H]; [1.61 - 1.75 (m); 1.91 - 2.05 (m); 1H]; [3.42 (d, J = 7.8 Hz), 3.60 (d, J = 6.4 Hz); 1H]; [3.75 - 3.85 (m), 3.94 - 4.04 (m), 1H]; [4.22 (s),
15 4.26 (s); 2H]; 4.95 - 5.10 (m, 3H); [5.24 (d, J = 9.4 Hz), 5.27 (d, J = 9.9 Hz), 1H]; 7.20 - 7.60 (m, 9H).

j. 1-[3-[5-(3-Trifluoromethylbenzyl)]-1,2,4-oxadiazolyl]-2-(S)-Amino-3-Methyl-Butan-1-ol
20 Hydrochloride.

The amine, 1-[3-[5-(3-trifluoromethylbenzyl)]-1,2,4-oxadiazolyl]-2-(S)-benzyloxycarbonylamino-3-methylbutan-1-ol, was dissolved into 20 mL trifluoroacetic acid at room temperature. Once
25 dissolved, the mixture was cooled in an ice bath and thioanisole was added. The resulting mixture was allowed to warm to room temperature overnight and evaporated to give an oil. The oil was dissolved into 50 mL of dry ether (0°C) and 1.5 equivalents of HCl in
30 dioxane was added to give a solid. The suspension was allowed to settle and the ether decanted. An additional volume of ether was added (50 mL) and decanted to remove remaining thioanisole. The resulting solid slowly became an oil which was dried
35 under vacuum (~ 1 mm Hg) for 5 h to give 0.96 g (49.4%) of 1-[3-[5-(3-trifluoromethylbenzyl)]-1,2,4-oxadiazolyl]-2-(S)-amino-3-methylbutan-1-ol hydrochloride.

¹H NMR (CDCl₃) δ 0.98 - 1.50 (m, 6H); 1.86 - 2.06 (m, 1H); 3.58 - 3.80 (m, 1H); 4.20 - 4.40 (m, 2H), [5.25 (d, J = 8.0 Hz), 5.37 (d, J = 3.4 Hz), 1H], 6.60 - 7.20 (brs, 1H), 7.20 - 7.70 (m, 4H), 7.90 - 8.20 (brs, 3H).

k. (Benzyloxycarbonyl)-L-Valyl-N-[1-[(3-[5-(3-Trifluoromethyl- benzyl)-1,2,4-oxadiazolyl]hydroxymethyl]-2-(S)-Methylpropyl]-L-Pro linamide.


A solution of Cbz-Val-Pro-OH (1.09 g, 2.83 mmol) in 30 mL of dry DMF was cooled in a brine/ice bath. To this stirred mixture was added HOBt (0.45 g, 3.34 mmol) followed by EDCI (0.52 g, 2.70 mmol). After stirring for 30 minutes, 1-[3-[5-(3-trifluoromethylbenzyl)]-1,2,4-oxadiazolyl]-2-(S)-amino-3-methyl-butan-1-ol hydrochloride (0.94 g, 2.57 mmol) in 20 mL of dry DMF was added dropwise followed by N-methyl morpholine (0.39 g, 3.85 mmol) and the reaction allowed to stir overnight at 0°C. The mixture was diluted with ethyl acetate (≈ 250 mL) and washed with saturated NaHCO₃ (2x), 5% KHSO₄, brine and the organics dried with anhydrous magnesium sulfate. The mixture was filtered and the solvent removed under reduced pressure. The residue was purified by column chromatography (silica gel, ethyl acetate:hexane 60:40 to 75:25) to give (benzyloxycarbonyl)-L-valyl-N-[1-[(3-[5-(3-trifluoromethylbenzyl)-1,2,4-oxadiazolyl]hydroxymethyl]-2-(S)-methylpropyl]-L-prolinamide as a white foamy solid; 0.88 g (51.9%).

¹H NMR (CDCl₃) δ 0.75 - 1.15 (m, 12H); 1.70 - 2.30 (m, 5.5H), 2.40 - 2.65 (m, 0.5H); 3.45 - 3.60 (m, 1H); 3.61 - 5.85 (m, 1H), 3.90 - 4.05 (m, 1H); 4.10 - 4.65 (m, 2H); [4.23 (s), 4.30 (s), 2H]; 4.75 - 5.15 (m, 3H); 5.30 - 5.85 (m, 1H); [6.91 (d, J = 9.2 Hz), 7.08 (d, J = 9.5 Hz), 1H]; 7.15 - 7.80 (m, 9H); [8.02 (d, J = 8.6 Hz), 8.38 (d, J = 8.6 Hz), 1H];

EXAMPLE II

(Benzyloxycarbonyl)-L-Valyl-N-[1-[(3-[5-(2-Phenylethyl)-1,2,4-oxadiazolyl]carbonyl]-2-(S)-Methylpropyl]-L-Prolinamide.

- 5 The compound was prepared using a similar oxidative procedure as described in Example I but utilizing (benzyloxycarbonyl)-L-valyl-N-[1-[(3-[5-(2-phenylethyl)-1,2,4-oxadiazolyl]hydroxymethyl]-2-(S)-methylpropyl]-L-prolinamide for the alcohol; an
- 10 analytical sample was obtained from RP-HPLC (isocratic, CH₃CN: H₂O: 3:2) as a white solid; ¹H NMR (CDCl₃) δ [0.89 (d, J = 6.9 Hz); 0.96 (d, J = 6.9 Hz); 1.02 (d, J = 6.9 Hz), 12H]; 1.80 - 2.20 (m, 4H); 2.22 - 2.40 (m, 2H); 3.18 - 3.36 (m, 5H); 3.57 - 3.69 (m, 1H); 3.70 - 3.86 (m, 1H); 4.36 (dd, J₁ = 6.6 Hz; J₂ = 9.0 Hz; 1H); 4.63 (dd, J₁ = 2.9 Hz, J₂ = 8.2 Hz, 1H); 5.10 (ABq, J = 12.4 Hz, 2H); 5.32 (dd, J₁ = 4.9 Hz, J₂ = 7.6 Hz, 1H); 5.66 (d, J = 9.1 Hz, 1H); 7.18 - 7.40 (m, 10H); 7.45 (d, J = 7.40 Hz, 1H)
- 15 ¹³C NMR (CDCl₃) δ 17.24, 17.59, 19.45, 19.76, 25.11, 27.23, 28.41, 30.41, 31.44, 32.39, 47.76, 57.52, 59.76, 61.49, 66.93, 126.82, 127.99, 128.09, 128.17, 128.48, 128.73, 136.32, 138.88, 156.44, 164.74, 171.09, 172.20, 180.62, 190.18
- 20 IR (Deposit) 3298.4, 2964.8, 1719.1, 1683.9, 1630.6, 1525.7, 1452.9, 1230.4 cm⁻¹.



	%C	%H	%N
30 Theory	65.65	6.85	11.60
Found	65.50	6.68	11.61

- The intermediate (benzyloxycarbonyl)-L-valyl-N-[1-[(3-[5-(2-phenylethyl)-1,2,4-oxadiazolyl]hydroxymethyl]-2-(S)-methylpropyl]-L-prolinamide was prepared as follows:
- 35

a. 3-Phenylpropionyl Chloride.

To a mixture of 3-phenyl propionic acid (15.00 g, 0.10 mol) in 50 mL of dry methylene chloride was added 10.9 mL (0.15 mol) of oxalyl chloride over 5 minutes.

- 5 The mixture was allowed to stir at RT overnight. The mixture was evaporated (40 °C, aspirator vacuum) to an oil which was distilled bp_{0.9mm Hg} 74-75 °C to afford 15.14 g (89.9%) of product as a clear colorless liquid.

10 IR (Neat) 1790.4cm⁻¹.

b. 1-[(N-Hydroxy)carboximid-N-(3-phenylpropionyl)amido]-1-Acetoxy-2-(S)-Benzyloxycarbonylamino-3-Methyl-Butane.

- 15 This compound was prepared following substantially the same procedure as described in Example I, except 3-phenylpropionyl chloride was used as the acid chloride; 49.1%, off-white solid.

¹H NMR (CDCl₃) δ [0.89 (d, J = 6.8 Hz); 0.93 (d, J = 6.8 Hz); 0.96 (d, J = 6.8 Hz); 1.00 (d, J = 6.8 Hz); 6H]; 1.82 - 1.96 (m, 1H); [1.91 (s), 2.06 (s), 3H]; 2.66 - 2.80 (m, 2H); 2.92 - 3.04 (m, 2H); 3.82 - 3.94 (m, 1H); 4.76 - 5.42 (m, 6H), 7.10 - 7.50 (m, 10H).

25

c. 1-[3-[5-(2-Phenethyl)]-1,2,4-oxadiazolyl]-1-Acetoxy-2-(S)-Benzyloxycarbonylamino-3-Methyl-Butane.

- This compound was prepared by a cyclization procedure as described in Example I utilizing 1-[(N-hydroxy)carboximid-N-(3-phenylpropionyl)amido]-1-acetoxy-2-(S)-benzyloxycarbonylamino-3-methyl-butane; oil (yield 86.7%).

¹H (CDCl₃) δ [0.90 (d, J = 6.8 Hz); 0.97 (d, J = 6.8 Hz); 0.98 (d, J = 6.8 Hz); 1.02 (d, J = 6.8 Hz); 6H]; [1.28 - 1.44 (m); 1.68 - 1.84 (m); 1H]; [2.05 (s), 2.12 (s); 3H]; 3.04 - 3.28 (m, 4H); 3.98 - 4.10 (m, 1H); [5.04 (s), 5.13 (s); 2H]; 5.02 - 5.06 (m,

35

1H); 5.41 (d, J = 10.1 Hz; 1H); [6.06 (d, J = 3.5 Hz); 6.09 (d, J = 7.8 Hz); 1H]; 7.12 - 7.46 (m, 10H).

- d. 1-[3-[5-(2-phenethyl)]-1,2,4-oxadiazolyl]-2-(S)-
5 Benzyloxycarbonylamino-3-Methyl-Butan-1-ol.

The compound was prepared using the hydrolysis conditions in Example I with 1-[3-[5-(2-phenethyl)]-1,2,4-oxadiazolyl]-1-acetoxy-2-(S)-benzyloxycarbonylamino-3-methyl-butane, oil (yield
10 76.7%).

¹H (CDCl₃) δ [0.91 (d, J = 6.9 Hz); 0.96 (d, J = 6.9 Hz); 0.99 (d, J = 6.9 Hz); 1.08 (d, J = 6.9 Hz); 6H]; [1.38 - 1.52 (m), 1.91 - 2.03 (m), 1H]; 3.02 - 3.26 (m, 5H); [3.76 - 3.86 (m), 3.88 - 4.00 (m) 1H];
15 4.92 - 5.28 (m, 4H), 7.15 - 7.50 (m, 10H).

- e. 1-[3-[5-(2-Phenethyl)]-1,2,4-oxadiazolyl]-2-(S)-
Amino-3-Methyl-Butan-1-ol Hydrochloride.

This compound was prepared following
20 substantially the same procedure as described in Example I utilizing 1-[3-[5-(2-phenethyl)]-1,2,4-oxadiazolyl]-2-(S)-benzyloxycarbonylamino-3-methylbutan-1-ol; Heavy oil (yield 81.3%); ¹H (CDCl₃) δ [1.03 (d, J = 6.9 Hz); 1.05 (d, J = 6.9 Hz); 1.09 (d, J = 6.9 Hz); 6H]; [1.76 - 1.87 (m), 1.87 - 2.01 (m), 1H]; 3.02 - 3.30 (m, 5H); [3.62 - 3.73 (m), 3.73 - 3.83 (m), 1H]; [5.28 (d, J = 9.0 Hz); 5.44 (d, J = 3.5 Hz); 1H]; 6.74 - 7.06 (brs, 1H); 7.10 - 7.42 (m, 5H); 8.00 - 8.20 (2brs, 3H).
30

- f. (Benzyloxycarbonyl)-L-Valyl-N-[1-[(3-[5-(2-Phenethyl)-1,2,4-oxadiazolyl]hydroxymethyl)-2-(S)-Methylpropyl]-L-Prolinamide.

This compound was prepared by a similar coupling
35 procedure as described in Example I using the hydrochloride of 1-[3-[5-(2-phenethyl)]-1,2,4-oxadiazolyl]-2-(S)-amino-3-methyl-butane-1-ol to give (benzyloxycarbonyl)-L-valyl-N-[1-[(3-[5-(2-

phenylethyl)-1,2,4-oxadiazolyl]hydroxymethyl]-2-(S)-methylpropyl]-L-prolinamide as a white solid (yield 46.8%).

^1H (CDCl_3) δ 0.70 - 1.14 (m, 12H); 1.50 - 2.60 (m, 6H); 2.90 - 3.30 (m, 3H); 3.50 - 4.90 (m, 5H); 4.90 - 5.30 (m, 9H); 5.40 - 5.75 (m, 1H); 6.75 - 7.65 (m, 10H).

EXAMPLE III


(Benzyloxycarbonyl)-L-Valyl-N-[1-[(3-[5-(2-Methoxybenzyl)-1,2,4-oxadiazolyl]carbonyl]-2-(S)-Methylpropyl]-L-Prolinamide.

The compound was prepared using a similar oxidative procedure as described in Example I but utilizing (benzyloxycarbonyl)-L-valyl-N-[1-[(3-[5-(2-methoxybenzyl)-1,2,4-oxadiazolyl]hydroxymethyl]-2-(S)-methylpropyl]-L-prolinamide for the alcohol; R_f = 0.74 (silica gel 4:1, ethyl acetate:hexane). An analytical sample was obtained from RP-HPLC (isocratic, CH_3CN : H_2O : 3:2) as a white solid; TLC, R_f = 0.74 ethylacetate:hexane (4:1).

^1H NMR (CDCl_3) δ [0.88 (d, J = 7.0 Hz); 0.95 (d, J = 7.0 Hz); 1.01 (d, J = 7.0 Hz), 12H]; 1.80 - 2.20 (m, 4H); 2.20 - 2.45 (m, 2H); 3.55 - 3.80 (m, 1H); 3.70 - 3.80 (m, 1H); 3.80 (s, 3 H), 4.35 (s, 2 H), 4.35 (dd, J_1 = 6.8 Hz; J_2 = 9.1 Hz; 1H); 4.62 (dd, J_1 = 2.6 Hz, J_2 = 7.8 Hz, 1H); 5.10 (ABq, J = 12.4 Hz, 2H); 5.33 (dd, J_1 = 5.0 Hz, J_2 = 7.6 Hz, 1H); 5.61 (d, J = 9.2 Hz, 1H); 6.87 - 7.00 (m, 2 H); 7.20 - 7.50 (m, 7H).

^{13}C NMR (CDCl_3) δ 17.12, 17.55, 19.47, 19.81, 25.12, 27.26, 27.82, 30.42, 31.41, 47.75, 55.41, 57.46, 59.77, 61.37, 66.93, 110.67, 120.74, 127.99, 128.10, 128.49, 129.40, 130.60, 136.27, 156.41, 157.22, 164.73, 171.03, 172.14, 180.17, 190.20.

IR (Deposit) 3305.5, 2965.4, 1716.6, 1682.7, 1633.5 cm^{-1} .



	%C	%H	%N
Theory	63.96	6.67	11.30
Found	63.78	6.49	11.19

5

The intermediate (benzyloxycarbonyl)-L-valyl-N-[1-[(3-[5-(2-methoxybenzyl)-1,2,4-oxadiazolyl]hydroxymethyl)-2-(S)-methylpropyl]-L-prolinamide was prepared as follows:

10

a. 2-Methoxyphenylacetyl Chloride.

To a mixture of 2-methoxyphenylacetic acid (10.00 g, 0.060 mol) in 40 mL of dry methylene chloride was added 7.88 mL (0.090 mol) of oxalyl chloride over 5 minutes. The mixture was allowed to proceed overnight. The mixture was evaporated (40°C, aspirator vacuum) to an oil which was used without further purification; 10.89 g (98.0 %) clear colorless liquid.

20 IR (Neat) 1803.5 cm^{-1} .

b. 1-[(N-Hydroxy)carboximid-N-(2-methoxybenzyl)amido]-1-Acetoxy-2-(S)-Benzyloxycarbonylamino-3-Methyl-Butane.

25 This compound was prepared following substantially the same procedure as described in Example I, but 2-methoxyphenylacetyl chloride using as the acid chloride; 32.1%, off-white solid.

^1H NMR (CDCl_3) δ [0.89 (d, $J = 7.0$ Hz); 0.93 (d, $J = 7.0$ Hz); 0.99 (d, $J = 7.0$ Hz); 1.01 (d, $J = 7.0$ Hz); 6H]; 1.82 - 2.20 (m, 1H); [1.91 (s), 2.06 (s), 3H]; [3.73 (s), 3.75 (s), 3.79 (s), 3.81 (s), 3H]; 3.80 - 3.98 (m, 1H); 4.84 - 5.46 (m, 6H), 6.82 - 6.88 (m, 2H); 7.90 - 7.40 (m, 9H).

35

c. 1-[3-[5-(2-methoxybenzyl)]-1,2,4-oxadiazolyl]-1-Acetoxy-2-(S)-Benzyloxycarbonylamino-3-Methyl-Butane.

This compound was prepared by a cyclization procedure as described in Example I utilizing 1-[(N-hydroxy)carboximid-N-(2-methoxybenzyl)amido]-1-acetoxy-2-(S)-benzyloxycarbonylamino-3-methyl-butane;
5 oil (yield 86.7%).

^1H (CDCl_3) δ [0.92 (d, J = 6.7 Hz); 0.97 (d, J = 6.7 Hz); 0.99 (d, J = 6.7 Hz); 1.02 (d, J = 6.7 Hz); 6H]; [1.45 - 1.60 (m); 1.70 - 1.82 (m); 1H]; [2.02 (s), 2.11 (s), 3H]; [3.76 (s), 3.77 (s), 3H]; 4.00 -
10 4.18 (m, 1H); [4.18 (s), 4.23 (s), 2H]; 4.99 - 5.14 (m, 2.5H); 5.59 (d, J = 10.0 Hz, 0.5H); [6.07 (d, J = 3.5 Hz); 6.12 (d, J = 4.7 Hz), 1H]; 6.84 - 6.98 (m, 2H); 7.16 - 7.40 (m, 7H).

15 d. 1-[3-[5-(2-methoxybenzyl)]-1,2,4-oxadiazolyl]-2-(S)-Benzyloxycarbonylamino-3-Methyl-Butan-1-ol.

This compound was prepared using the hydrolysis conditions in Example I but with the intermediate 1-[3-[5-(2-methoxybenzyl)]-1,2,4-oxadiazolyl]-1-acetoxy-
20 2-(S)-benzyloxycarbonylamino-3-methyl-butane, oil (yield 76.7%).

^1H (CDCl_3) δ [0.92 (d, J = 6.7 Hz); 0.95 (d, J = 6.7 Hz); 0.97 (d, J = 6.7 Hz); 1.05 (d, J = 6.7 Hz); 6H]; [1.52 - 1.66 (m); 1.90 - 2.30 (m); 1H]; [3.12 (d, J = 8.2 Hz), 3.43 (d, J = 5.9 Hz), 1H]; [3.75 (s),
25 3.76 (s), 3H]; [3.76 - 3.86 (m), 3.90 - 4.00 (m), 1H]; [4.18 (s), 4.20 (s); 2H]; 4.96 - 5.05 (m), 5.09 (ABq, J = 12.1 Hz, 2H); [5.19 (d, J = 9.8 Hz); 5.40 (d, J = 10.7 Hz); 1H]; 6.80 - 7.00 (m, 2H); 7.15 - 7.45 (m,
30 7H).

e. 1-[3-[5-(2-methoxybenzyl)]-1,2,4-oxadiazolyl]-2-(S)-Amino-3-Methyl-Butan-1-ol Hydrochloride.

This compound was prepared following
35 substantially the same procedure as described in Example I utilizing 1-[3-[5-(2-methoxybenzyl)]-1,2,4-oxadiazolyl]-2-(S)-benzyloxycarbonylamino-3-methyl-butan-1-ol; Heavy oil (yield 81.3%).

^1H (CDCl_3) δ [0.97 (d, $J = 7.0$ Hz); 1.00 (d, $J = 7.0$ Hz); 1.04 (d, $J = 7.0$ Hz); 6H]; 1.84 - 2.00 (m, 1H); 3.58 - 3.68 (m, 1H); [3.71 (s), 3.72 (s), 3H]; [4.16 (s), 4.20 (s), 2H]; [5.20 (d, $J = 8.0$ Hz), 5.30 (d, $J = 3.4$ Hz), 1H]; 6.80 - 6.92 (m, 2H); 7.10 - 7.45 (m, 2H); 7.90 - 8.20 (2brs, 3H).

f. (Benzyloxycarbonyl)-L-Valyl-N-[1-[(3-[5-(2-methoxybenzyl)-1, 2,4-oxadiazolyl]hydroxymethyl)-2-(S)-Methylpropyl]-L-Prolinamide.

This compound was prepared by a similar coupling procedure as described in Example I using the hydrochloride of 1-[3-[5-(2-methoxybenzyl)]-1,2,4-oxadiazolyl]-2-(S)-amino-3-methyl-butan-1-ol to give (benzyloxycarbonyl)-L-valyl-N-[1-[(3-[5-(2-phenylethyl)oxadiazolyl]hydroxymethyl)-2-(S)-methylpropyl]-L-prolinamide as a white solid (yield 46.8%).

^1H (CDCl_3) δ 0.80 - 1.60 (m, 12H); 1.60 - 2.20 (m, 6H); 3.50 - 4.70 (m, 12H); 4.95 - 5.25 (m, 3H); 6.80 - 7.00 (m, 2H), 7.10 - 7.50 (m, 7H).

EXAMPLE IV

(Benzyloxycarbonyl)-L-Valyl-N-[1-[(3-[5-(Trifluoromethyl)-1, 2,4-oxadiazolyl]carbonyl)-2-(S)-Methylpropyl]-L-Prolinamide.

The compound was prepared using a similar oxidative procedure as described in Example I but utilizing (benzyloxycarbonyl)-L-valyl-N-[1-[(3-[5-(trifluoromethyl)-1,2,4-oxadiazolyl]hydroxymethyl)-2-(S)-methylpropyl]-L-prolinamide for the alcohol; $R_f = 0.64$ (silica gel 1:1, ethyl acetate:hexane). Purification was conducted using column chromatography (silica gel 1:1, ethyl acetate:hexane) to give a white solid (51.7 % Yield).

^1H NMR (CDCl_3) δ 0.82-0.87 (2d overlapping, 6H); 0.90 (d, $J = 6.9$ Hz, 3H); 0.96 (d, $J = 6.9$ Hz, 3H); 1.60 - 2.10 (m, 5H); 2.25-2.45 (m, 1H); 3.45-3.60 (m,

1H); 3.65 - 3.80 (m, 1H); 4.00 (t, J = 8.3 Hz, 1H),
4.43 (dd, J₁ = 4.8 Hz, J₂ = 8.3 Hz, 1H), 4.81 (t, J =
6.0 Hz; 1H); 5.01 (ABq, J = 13.1 Hz, 2H); 7.25-7.45
(bs, 6H); 8.62 (d, J = 6.5 Hz, 1H).

5 ¹³C NMR (CDCl₃) δ 17.51, 18.41, 18.88, 19.56,
24.45, 28.60, 29.01, 29.74, 47.03, 57.81, 58.61,
61.71, 65.37, 127.62, 127.75, 128.31, 137.06, 156.16,
164.32, 170.04, 172.17, 188.17.

IR (Deposit) 3431, 3296, 3024, 2970, 2878, 1717,
10 1684, 1630 cm⁻¹.

C₂₆H₃₂N₅O₆F₃

	%C	%H	%N
Theory	55.02	5.68	12.34
15 Found	54.76	5.55	12.18

The intermediate (benzyloxycarbonyl)-L-valyl-N-[1-
[(3-[5-(trifluoromethyl)-1,2,4-
oxadiazolyl]hydroxymethyl]-2-(S)-methylpropyl]-L-
20 prolinamide was prepared as follows:

a. 1-[3-[5-(Trifluoromethyl)]-1,2,4-oxadiazolyl]-1-
Acetoxy-2-(S)-Benzyloxycarbonylamino-3-Methyl-Butane.

A solution containing 10.0 g (29.64 mmol) of 1-
25 [(N-hydroxy)carboximidamide]-1-acetoxy-2-(S)-
benzyloxycarbonylamino-3-methyl-butane and 12.4 mL
(87.79 mmol) of trifluoroacetic anhydride in 180 mL of
anhydrous toluene was heated to 90 °C under a nitrogen
atmosphere. The reaction was monitored by TLC (silica
30 gel; ethyl acetate:hexane 1:1). After 20 min, the
solvent was removed under reduced pressure and the
residue purified by column chromatography (silica gel;
ethyl acetate:hexane, 1:1) to afford 6.35 g (51.6 %) of
product as a viscous pale yellow oil.

35 ¹H (CDCl₃) δ 0.95-1.10 (m, 6H); 1.70 - 1.90 (m;
1H); [2.09 (s), 2.16 (s), 3H]; [3.94 - 4.06 (m), 4.08
- 4.20 (m) 1H]; [5.01 (s), 5.08 (s); 2H]; 4.95 - 5.15

(m, 1H); [6.08 (d, J = 6.2 Hz); 6.13 (d, J = 2.7 Hz); 1H]; 7.20 - 7.50 (m, 5H).

- b. 1-[3-[5-(Trifluoromethyl)]-1,2,4-oxadiazolyl]-2-
5 (S)-Benzyloxycarbonylamino-3-Methyl-Butan-1-ol.

This compound was prepared using hydrolysis conditions in Example I but with the intermediate 1-[3-[5-(trifluoromethyl)]-1,2,4-oxadiazolyl]-1-acetoxy-2-(S)-benzyloxycarbonylamino-3-methyl-butane, oil
10 (yield 72.7 %).

¹H (CDCl₃) δ [0.97 (d, J = 6.9 Hz), 1.01 (d, J = 6.9 Hz), 1.09 (d, J = 6.9 Hz), 6H]; [3.63 (d, J = 8.0 Hz), 3.73 (d, J = 7.1 Hz, 1H); [3.66 - 3.76 (m), 3.93 - 4.03 (m), 1H]; 4.95 - 5.24 (m, 3H), 7.26 - 7.45 (m,
15 5H).

- e. 1-[3-[5-(Trifluoromethyl)]-1,2,4-oxadiazolyl]-2-
(S)-Amino-3-Methyl-Butan-1-ol Hydrochloride.

This compound was prepared following
20 substantially the same procedure as described in Example I utilizing 1-[3-[5-(trifluoromethyl)]-1,2,4-oxadiazolyl]-2-(S)-benzyloxycarbonylamino-3-methylbutan-1-ol; Heavy oil (yield 83.8 %);

¹H (DMSO-d₆) δ 0.80 - 1.15 (m, 6H), 1.85 - 2.15
25 (m, 1H); 3.25 - 3.35 (m, 1H); 5.10 - 5.25 (brs, 1H); [8.10 (brs) 8.15 (brs), 3H].

- f. (Benzyloxycarbonyl)-L-Valyl-N-[1-[(3-[5-
(trifluoromethyl)-1, 2,4-oxadiazolyl]hydroxymethyl)-2-
30 (S)-Methylpropyl]-L-Prolinamide.

This compound was prepared by a similar coupling procedure as described in Example I using the hydrochloride of 1-[3-[5-(trifluoromethyl)]-1,2,4-oxadiazolyl]-2-(S)-amino-3-methyl-butan-1-ol to give
35 (benzyloxycarbonyl)-L-valyl-N-[1-[(3-[5-(trifluoromethyl)-1,2,4-oxadiazolyl]hydroxymethyl)-2-(S)-methylpropyl]-L-prolinamide as a white solid (yield 63.1 %); R_f = 0.54, 3:2 ethyl acetate:hexane.

^1H (CDCl_3) δ 0.80 - 1.20 (m, 6H); 1.80 - 2.60 (m, 6H); 3.50 - 4.70 (m, 6H); 4.90-5.30 (m, 3H); 5.40 - 5.75 (m, 1H); 6.85 - 7.20 (m, 1H), 7.25 - 7.40 (m, 5H).

5

EXAMPLE V

(Benzyloxycarbonyl)-L-Valyl-N-[1-[(3-[5-(methyl)-1,2,4-oxadiazolyl]carbonyl]-2-(S)-Methylpropyl]-L-Prolinamide.

10 The compound was prepared using a similar oxidative procedure as described in Example I but utilizing (benzyloxycarbonyl)-L-valyl-N-[1-[(3-[5-(methyl)-1,2,4-oxadiazolyl]hydroxymethyl]-2-(S)-methylpropyl]-L-prolinamide for the alcohol; R_f = 0.42
 15 (silica gel, 5:95, methanol:methylene chloride). Purification was conducted by column chromatography on silica gel (5:95, methanol:methylene chloride) to give a white solid (57.2%).

^1H NMR ($\text{DMSO}-d_6$) δ [0.90 (d, J = 6.9 Hz); 0.96
 20 (d, J = 6.9 Hz); 1.02 (d, J = 6.9 Hz), 12H]; 1.80 - 2.25 (m, 4H); 2.25 - 2.45 (m, 2H); 2.68 (s, 3H); 3.55 (dd, J_1 = 5.6 Hz, J_2 = 15.8 Hz, 1H); 3.71 (dd, J_1 = 7.1 Hz, J_2 = 16.0 Hz, 1H); 4.01 (t, J = 8.3 Hz, 1H); 4.47 (dd, J_1 = 4.3 Hz, J_2 = 7.9 Hz, 1H); 4.92 (t, J = 6.1
 25 Hz, 1H); 5.01 (ABq, J = 12.6 Hz, 2H); 7.35 (s, 5H); 7.40 (d, J = 7.0 Hz, 1H); 7.44 (d, J = 8.0 Hz, 1H).

^{13}C NMR (CDCl_3) δ 12.12, 17.14, 17.56, 19.26,
 19.62, 24.95, 27.22, 30.26, 31.27, 47.62, 57.44,
 59.52, 61.27, 66.72, 127.88, 127.94, 128.35, 136.21,
 30 156.37, 164.74, 171.13, 171.99, 178.01, 190.07.

$\text{C}_{26}\text{H}_{35}\text{N}_5\text{O}_6 \cdot 0.5 \text{H}_2\text{O}$

	%C	%H	%N
Theory	59.76	6.94	13.40
35 Found	59.98	6.69	13.16

The intermediate (benzyloxycarbonyl)-L-valyl-N-[1-[(3-[5-(methyl)-1,2,4-oxadiazolyl]hydroxymethyl]-2-

(S)-methylpropyl]-L-prolinamide was prepared as follows:

- a. 1-[3-[5-(Methyl)]-1,2,4-oxadiazolyl]-1-Acetoxy-2-
5 (S)-Benzyloxycarbonylamino-3-Methyl-Butane.

This compound was prepared using a similar cyclization procedure as reported in Example IV except acetic anhydride was used, oil (66.9 % yield).

¹H (CDCl₃) δ 0.94 (d, J = 6.7 Hz), 0.99 (d, J =
10 6.7 Hz), 1.00 (d, J = 6.7 Hz) 1.03 (d, J = 6.7 Hz),
6H); [1.44 - 1.66 (m), 1.72 - 1.86 (m), 1H]; [2.11
(s), 2.12 (s), 3H]; [2.53 (s), 2.60 (s), 3H]; 3.95 -
4.15 (m, 1H), 5.04 (ABq, J = 12.2 Hz, 2H), 5.04 - 5.14
(m, 1H); [6.05 (d, J = 3.4 Hz); 6.11 (d, J = 4.9 Hz),
15 1H]; 7.28 - 7.40 (m, 5H).

- b. 1-[3-[5-(Methyl)]-1,2,4-oxadiazolyl]-2-(S)-
Benzyloxycarbonylamino-3-Methyl-Butan-1-ol.

This compound was prepared using hydrolysis
20 conditions in Example I but with the intermediate 1-
[3-[5-(methyl)]-1,2,4-oxadiazolyl]-1-acetoxy-2-(S)-
benzyloxycarbonylamino-3-methyl-butane, oil (yield
93.2 %).

¹H (CDCl₃) δ [0.94 (d, J = 6.7 Hz), 0.98 (d, J =
25 6.7 Hz), 0.99 (d, J = 6.7 Hz), 1.07 (d, J = 6.7 Hz),
6H); [1.54 - 1.66 (m), 1.92 - 2.06 (m), 1H]; [2.54
(s), 2.58(s), 3H]; [3.09 (d, J = 8.2 Hz), 3.31 (d, J =
5.8 Hz), 1H]; [3.72 - 3.86 (m), 3.90 - 4.01 (m),
1H]; 4.97 - 5.20 (m, 3.5H), 5.35 (brd, J = 9.9 Hz,
30 0.5H); 7.25 - 7.45 (m, 5H).

- e. 1-[3-[5-(Methyl)]-1,2,4-oxadiazolyl]-2-(S)-Amino-
3-Methyl-Butan-1-ol Hydrochloride.

This compound was prepared following
35 substantially the same procedure as described in
Example I utilizing 1-[3-[5-(methyl)]-1,2,4-
oxadiazolyl]-2-(S)-benzyloxycarbonylamino-3-methyl-
butan-1-ol; Heavy oil (yield 87.3 %);

^1H (DMSO- d_6) δ [0.89 (d, J = 7.0 Hz); 0.95 (d, J = 7.0 Hz); 0.98 (d, J = 6.7 Hz); 6H]; 1.70 - 1.86 (m, 1H); 2.62 (s, 3H); 3.15 - 3.35 (m, 1H); 4.85-5.05 (m, 1H); 7.90 - 8.10 (brs, 3H).

5

f. (Benzyloxycarbonyl)-L-Valyl-N-[1-[(3-[5-(Methyl)-1,2,4-oxadiazolyl]hydroxymethyl)-2-(S)-Methylpropyl]-L-Prolinamide.

This compound was prepared by a similar coupling procedure as described in Example I using the hydrochloride of 1-[3-[5-(methyl)-1,2,4-oxadiazolyl]-2-(S)-amino-3-methyl-butan-1-ol to give (benzyloxycarbonyl)-L-valyl-N-[1-[(3-[5-(trifluoromethyl)oxadiazolyl]hydroxymethyl)-2-(S)-methylpropyl]-L-prolinamide as a white solid (yield 44.9 %); R_f = 0.74, 9:1 methylene chloride:methanol.

^1H (CDCl_3) δ 0.80 - 1.20 (m, 12H); 1.70 - 2.25 (m, 6H); [2.52 (s), 2.59 (s), 3H]; 3.50 - 3.85 (m, 2H); 3.90-4.45 (m, 3H); 4.45 - 4.65 (m, 1H); [4.69 (d, J = 8.1 Hz), 4.87 (d, J = 11.9 Hz), 1H]; 4.96 - 5.16 (m, 2H); [5.64 (d, J = 9.0 Hz), 5.85 (d, J = 9.0 Hz), 1H]; [6.91 (d, J = 9.2 Hz), 7.14 (d, J = 9.3 Hz), 1H], 7.34 (brs, 5H).

25 EXAMPLE VI

(Benzyloxycarbonyl)-L-Valyl-N-[1-[(3-[5-(Difluoromethyl)-1,2,4-oxadiazolyl]carbonyl)-2-(S)-Methylpropyl]-L-Prolinamide.


The compound was prepared using a similar oxidative procedure as described in Example I but utilizing (benzyloxycarbonyl)-L-valyl-N-[1-[(3-[5-(difluoromethyl)-1,2,4-oxadiazolyl]hydroxymethyl)-2-(S)-methylpropyl]-L-prolinamide for the alcohol; R_f = 0.48 (silica gel, 3:2, ethyl acetate:hexane).

Purification was conducted by column chromatography on silica gel (1:1, ethyl acetate:hexane) to give a white solid (yield 49.2%).

¹H NMR (CDCl₃) δ 0.93 (d, J = 7.0 Hz, 3H); 0.96 (d, J = 7.0 Hz, 3H); 1.01 (d, J = 7.0 Hz, 3H); 1.03 (d, J = 7.0 Hz), 1.80 - 2.20 (m, 4H); 2.25-2.45 (m, 2H); 3.55-3.65 (m, 1H), 3.65 - 3.85 (m, 1H), 4.35 (t, J = 7.0 Hz, 1H); 4.64 (d, J = 6.0 Hz, 1H); 5.01 (ABq, J = 12.2 Hz, 2H); 5.21 (t, J = 5.6 Hz, 1H), 5.52 (d, J = 9.4 Hz, 1H), 6.88 (t, J = 51.7 Hz, 1H), 7.36 (s, 5H), 7.62 (d, J = 6.5 Hz, 1H).

¹³C NMR (CDCl₃) δ 17.54, 17.63, 19.47, 19.77, 25.16, 26.81, 30.07, 31.46, 47.85, 57.54, 59.59, 61.96, 67.03, 105.31 (t, J = 195 Hz), 128.04, 128.18, 128.54, 136.27, 156.42, 164.71, 171.17, 172.53, 188.78.

IR (Deposit) 3430, 3304, 3027, 2969, 2878, 1717, 1682, 1630 cm⁻¹.



	%C	%H	%N
Theory	56.82	6.05	12.74
Found	56.81	5.97	12.62

The intermediate (benzyloxycarbonyl)-L-valyl-N-[1-[(3-[5-(difluoromethyl)-1,2,4-oxadiazolyl]hydroxymethyl)-2-(S)-methylpropyl]-L-prolinamide was prepared as follows:

a. 1-[3-[5-(Dimethyl)]-1,2,4-oxadiazolyl]-1-Acetoxy-2-(S)-Benzyloxycarbonylamino-3-Methyl-Butane.

This compound was prepared using a similar cyclization procedure as reported in Example IV except difluoroacetic anhydride was used, oil (36.0 % yield).

¹H (CDCl₃) δ [0.94 (d, J = 6.7 Hz), 0.99 (d, J = 6.7 Hz), 1.01 (d, J = 6.7 Hz), 6H], 1.04 (d, J = 6.7 Hz), 6H]; 1.58 - 1.92 (m; 1H), [2.10 (s), 2.15 (s), 3H]; 3.90 - 4.20 (m, 1H); 4.95 - 5.25 (m, 3 H), [5.94 (t, J = 53.2 Hz, 5.97 (t, J = 53.2 Hz), 1H], [6.12 (d, J = 5.2 Hz, 6.25 (d, J = 5.0 Hz), 1H); 7.30 - 7.45 (m, 5H).

b. 1-[3-[5-(Difluoromethyl)]-1,2,4-oxadiazolyl]-2-(S)-Benzyloxycarbonylamino-3-Methyl-Butan-1-ol.

This compound was prepared using hydolysis conditions in Example I but with the intermediate 1-
5 [3-[5-(difluoromethyl)]-1,2,4-oxadiazolyl]-1-acetoxy-2-(S)-benzyloxycarbonylamino-3-methyl-butane, oil (yield 70.1 %).

^1H (CDCl_3) δ [0.95 (d, $J = 6.7$ Hz), 1.00 (d, $J = 6.7$ Hz), 1.08 (d, $J = 6.7$ Hz), 6H]; [1.62 - 1.86 (m),
10 1.98 - 2.14 (m), 1H]; [3.25 (d, $J = 6.6$ Hz), 3.60 (d, $J = 6.7$ Hz), 1H]; [3.97 (ddd, $J_1 = 5.3$ Hz, $J_2 = 7.9$ Hz, $J_3 = 9.8$ Hz), 4.31 (ddd, $J_1 = 5.5$ Hz, $J_2 = 7.8$ Hz, $J_3 = 9.7$ Hz), 1H]; [5.03 (s), 5.10 (s), 2H]; 4.98 - 5.22 (m, 2H); [5.94 (t, $J = 54.3$ Hz), 6.72 (t, $J =$
15 54.3 Hz), 1H]; 7.27 - 7.40 (m, 5H).

e. 1-[3-[5-(Difluoromethyl)]-1,2,4-oxadiazolyl]-2-(S)-Amino-3-Methyl-Butan-1-ol Hydrochloride.

This compound was prepared following
20 substantially the same procedure as described in Example I utilizing 1-[3-[5-(difluoromethyl)]-1,2,4-oxadiazolyl]-2-(S)-benzyloxycarbonylamino-3-methyl-butan-1-ol; heavy oil (yield 81.8 %);

^1H ($\text{DMSO}-d_6$) δ [0.93 (d, $J = 7.0$ Hz); 0.96 (d, $J = 7.0$ Hz); 0.98 (d, $J = 7.0$ Hz); 1.01 (d, $J = 7.0$ Hz),
25 6H]; 1.80 - 2.00 (m, 1H); 3.20 - 3.35 (m, 1H); [5.08 (t, $J = 5.9$ Hz), 5.15 (t, $J = 4.8$ Hz), 1H]; [7.30 (t, $J = 53$ Hz), 7.55 (t, $J = 53$ Hz), 1H], 8.05 (brs, 3H).

30 f. (Benzyloxycarbonyl)-L-Valyl-N-[1-[(3-[5-(Difluoromethyl)]-1,2,4-oxadiazolyl]hydroxymethyl)]-2-(S)-Methylpropyl]-L-Prolinamide.

This compound was prepared by a similar coupling procedure as described in Example I using the
35 hydrochloride of 1-[3-[5-(difluoromethyl)]-1,2,4-oxadiazolyl]-2-(S)-amino-3-methyl-butan-1-ol to give (benzyloxycarbonyl)-L-valyl-N-[1-[(3-[5-(difluoromethyl)]-1,2,4-oxadiazolyl]hydroxymethyl)]-2-

(S)-methylpropyl]-L-prolinamide as a white solid
(yield 47.1 %); R_f = 0.27, 65% ethyl acetate:hexane.

^1H (CDCl_3) δ 0.84 - 1.18 (m, 12H); [1.78 - 2.20
(m), 2.20 - 2.32 (m), 6H]; 3.50 - 3.65 (m, 1H); 3.66-
5 3.92 (m, 1H); 4.11 - 4.71 (m, 4H); 5.00 - 5.15 (m,
2.5H); 5.17 (dd, J_1 = 7.4 Hz, J_2 = 7.8 Hz, 0.5H);
[5.49 (bd, J = 9.3 Hz), 5.64 (d, J = 8.8 Hz), 1H];
[6.77 (t, J = 52.1 Hz), 6.80 (t, J = 52.1 Hz), 1H];
[6.88 (d, J = 8.7 Hz), 7.12 (d, J = 8.5 Hz), 1H]; 7.30
10 -7.45 (m, 5H).

EXAMPLE VII


(Benzyloxycarbonyl)-L-Valyl-N-[1-[(3-[5-(benzyl)-
1,2,4-oxadi azolyl]carbonyl]-2-(S)-Methylpropyl]-L-
15 Prolinamide.

The compound was prepared using a similar
oxidative procedure as described in Example I but
utilizing (benzyloxycarbonyl)-L-valyl-N-[1-[(3-[5-
(benzyl)-1,2,4-oxadiazolyl]hydroxymethyl]-2-(S)-
20 methylpropyl]-L-prolinamide for the alcohol; R_f = 0.30
(silica gel 3:2, ethyl acetate:hexane). The material
was purified by column chromatography on silica gel
(3:2, ethyl acetate:hexane) to give a white solid,
59.7 %yield.

25 ^1H NMR (CDCl_3) δ [0.88 (d, J = 6.8 Hz); 0.94 (d,
 J = 6.8 Hz); 1.00 (d, J = 6.8 Hz), 12H]; 1.80 - 2.20
(m, 4H); 2.20-2.45 (m, 2H); 3.55 - 3.70 (m, 1H); 3.75
- 3.85 (m, 1H); 4.30 (s, 2H), 4.30 - 4.40 (m, 1H),
4.61 (dd, J_1 = 2.4 Hz; J_2 = 7.7 Hz; 1H); 5.09 (ABq, J
30 = 12.0 Hz, 2H); 5.30 (dd, J_1 = 5.1 Hz, J_2 = 7.1 Hz,
1H); 5.60 (d, J = 9.3 Hz, 1H); 7.25 - 7.50 (m, 11H).

^{13}C NMR (CDCl_3) δ 17.07, 17.59, 19.29, 19.69,
24.97, 27.26, 30.19, 31.27, 32.82, 47.64, 57.46,
59.56, 61.31, 66.75, 127.75, 127.90, 127.96, 128.37,
35 128.86, 128.91, 132.47, 136.24, 156.39, 164.71,
171.14, 172.02, 179.44, 189.97.

IR (Deposit) 3429, 3027, 3013, 2969, 1721, 1683,
1635, 1574, 1509, 1436 cm^{-1} .



	%C	%H	%N
Theory	64.20	6.73	11.70
Found	64.00	6.77	11.62

5

The intermediate (benzyloxycarbonyl)-L-valyl-N-[1-[(3-[5-(benzyl)-1,2,4-oxadiazolyl]hydroxymethyl]-2-(S)-methylpropyl]-L-prolinamide was prepared as follows:

10

a. 1-[3-[5-(Benzyl)]-1,2,4-oxadiazolyl]-1-Acetoxy-2-(S)-Benzyloxy carbonylamino-3-Methyl-Butane.

This compound was prepared using a similar cyclization procedure as reported in Example IV except
15 phenylacetic anhydride was used, the oil was used without further purification; TLC R_f = 0.77, silica gel, 3:2 (ethyl acetate:hexane).

b. 1-[3-[5-(Benzyl)]-1,2,4-oxadiazolyl]-2-(S)-
20 Benzyloxycarbonyl amino-3-Methyl-Butan-1-ol.

This compound was prepared using the hydrolysis conditions in Example I but with the intermediate 1-[3-[5-(benzyl)]-1,2,4-oxadiazolyl]-1-acetoxy-2-(S)-benzyloxycarbonylamino-3-methyl-butane, oil (yield
25 47.0%); TLC R_f = 0.47, silica gel, 1:1 (ethyl acetate:hexane).

^1H (CDCl_3) δ [0.92 (d, J = 6.7 Hz), 0.98 (d, J = 6.7 Hz), 1.06 (d, J = 6.7 Hz), 6H]; [1.56 - 1.72 (m); 1.92 - 2.20 (m); 1H]; [3.00 - 3.08 (bd), 3.22 - 3.30
30 (m), 1H]; 3.74 - 3.86 (m, 0.5 H); 3.90 - 4.00 (m, 0.5H); [4.18 (s), 4.21 (s); 2H]; 4.94 - 5.32 (m, 4H); 7.20 - 7.50 (m, 10H).

c. 1-[3-[5-(Benzyl)]-1,2,4-oxadiazolyl]-2-(S)-Amino-
35 3-Methyl-Butan-1-ol Hydrochloride.

This compound was prepared following substantially the same procedure as described in Example I utilizing 1-[3-[5-(benzyl)]-1,2,4-

oxadiazolyl]-2-(S)-benzyloxycarbonylamino-3-methylbutan-1-ol; Heavy oil (yield 74.9%).

¹H (CDCl₃) δ [0.99 (d, J = 7.0 Hz); 1.01 (d, J = 7.0 Hz); 1.06 (d, J = 7.0 Hz); 1.07 (d, J = 7.0 Hz);
5 6H]; 1.80 - 2.10 (m, 1H); 3.60 - 3.80 (m, 1H); 4.10 - 4.28 (m, 2H); [5.24 (d, J = 8.3 Hz), 5.35 (d, J = 2.6 Hz), 1H]; 6.10 - 6.60 (bs, 1H); 7.10 - 7.45 (m, 5H); 8.17 (brs, 3H).

10 f. (Benzyloxycarbonyl)-L-Valyl-N-[1-[(3-[5-(Benzyl)-1,2,4-oxadiazolyl]hydroxymethyl)-2-(S)-Methylpropyl]-L-Prolinamide.

This compound was prepared by a similar coupling procedure as described in Example I using the
15 hydrochloride of 1-[3-[5-(benzyl)]-1,2,4-oxadiazolyl]-2-(S)-amino-3-methylbutan-1-ol to give (benzyloxycarbonyl)-L-valyl-N-[1-[(3-[5-(benzyl)-1,2,4-oxadiazolyl]hydroxymethyl)-2-(S)-methylpropyl]-L-prolinamide as a white solid (yield 74.7%); TLC R_f
20 = 0.10, silica gel, 3:2 (ethyl acetate:hexane).

¹H (CDCl₃) δ 0.70 - 1.20 (m, 12H); [1.65 - 2.20 (m), 2.45 - 2.60 (m), 6H]; 3.42 - 3.60 (m, 1H); 3.62 - 3.75 (m, 1H); 3.85 - 4.02 (m, 1H); 4.10 - 4.65 (m, 4H); 4.71 - 5.15 (m, 3H); 5.30 - 5.65 (m, 1H); [5.79
25 (d, J = 9.2 Hz), 6.85 (d, J = 8.5 Hz), 1H]; [7.03 (d, J = 9.8 Hz), 7.70 (brd), 1H]; 7.20 - 7.45 (m, 10H).

EXAMPLE VIII

(Benzyloxycarbonyl)-L-Valyl-N-[1-[(3-[5-(3-methoxybenzyl)-1,2,4-oxadiazolyl]carbonyl)-2-(S)-Methylpropyl]-L-Prolinamide.

The compound was prepared using a similar oxidative procedure as described in Example I but utilizing (benzyloxycarbonyl)-L-valyl-N-[1-[(3-[5-(3-methoxybenzyl)-1,2,4-oxadiazolyl]hydroxymethyl)-2-(S)-methylpropyl]-L-prolinamide for the alcohol; R_f = 0.60
35 (silica gel 4:1, ethyl acetate:hexane). An

analytical sample was obtained from RP-HPLC (isocratic, CH₃CN: H₂O: 60:40) as a white solid.

¹H NMR (CDCl₃) δ [0.88 (d, J = 6.9 Hz); 0.96 (d, J = 6.9 Hz); 1.02 (d, J = 6.9 Hz), 12H]; 1.85 - 2.20 (m, 4H); 2.20-2.40 (m, 2H); 3.55-3.70 (m, 1H); 3.70 - 3.85 (m, 1H); 3.80 (s, 3 H), 4.28 (s, 2H), 4.35 (dd, J₁ = 6.9 Hz; J₂ = 8.8 Hz; 1H); 4.62 (dd, J₁ = 2.6 Hz, J₂ = 7.8 Hz, 1H); 5.10 (ABq, J = 12.4 Hz, 2H); 5.31 (dd, J₁ = 4.9 Hz, J₂ = 7.5 Hz, 1H); 5.59 (d, J = 9.0 Hz, 1H); 6.80 - 7.00 (m, 3H); 7.20 - 7.45 (m, 7H).

¹³C NMR (CDCl₃) δ 17.14, 17.60, 19.46, 19.84, 25.12, 27.28, 30.36, 31.38, 32.99, 47.83, 55.28, 57.34, 59.85, 61.48, 66.99, 113.43, 114.66, 121.25, 128.00, 128.14, 128.51, 130.09, 133.88, 136.25, 156.45, 160.00, 164.76, 171.07, 172.35, 179.49, 189.99.

IR (Deposit) 3303.2, 2965.8, 1720.2, 1682.0, 1632.0 cm⁻¹.

20 C₃₃H₄₁N₅O₇ · 0.5 H₂O

	%C	%H	%N
Theory	63.04	6.73	11.14
Found	63.05	6.89	11.06

25 The intermediate (benzyloxycarbonyl)-L-valyl-N-[1-[(3-[5-(3-methoxybenzyl)-1,2,4-oxadiazolyl]hydroxymethyl]-2-(S)-methylpropyl]-L-prolinamide was prepared as follows:

30 a. 3-Methoxyphenylacetyl Chloride.

This acid chloride was prepared in a similar manner as for the acid chloride in Example I and was used without further purification.

35 b. 1-[(N-Hydroxy)carboximid-N-(3-methoxybenzyl)amido]-1-Acetoxy -2-(S)-Benzyloxycarbonylamino-3-Methyl-Butane.

This compound was prepared following substantially the same procedure as described in Example I, but 3-methoxyphenylacetyl chloride using as the acid chloride; 65.3 %, off-white solid.

5 ¹H NMR (CDCl₃) δ [0.89 (d, J = 6.9 Hz); 0.93 (d, J = 6.9 Hz); 0.96 (d, J = 6.9 Hz); 1.00 (d, J = 6.9 Hz); 6H]; 1.80 - 2.15 (m, 1H); [1.91 (s), 2.06 (s), 3H]; 3.60 - 3.95 (m, 1H); [3.70 (s), 3.74 (s), 3.78 (s), 3.79 (s), 3H]; 4.70 - 5.30 (m, 7H), [5.38 (d, J =
10 7.0 Hz), 5.41 (d, J = 6.3 Hz), 1H]; 6.78 - 6.92 (m, 3H); 7.18 - 7.40 (m, 6H).

c. 1-[3-[5-(3-Methoxybenzyl)]-1,2,4-oxadiazolyl]-1-Acetoxy-2-(S)-Benzyloxycarbonylamino-3-Methyl-Butane.

15 This compound was prepared in a similar manner by a cyclization procedure described in Example I utilizing 1-[(N-hydroxy)carboximid-N-(3-methoxybenzyl)amido]-1-acetoxy-2-(S)-benzyloxycarbonylamino-3-methyl-butane; oil (yield
20 77.2 %).

¹H (CDCl₃) δ [0.93 (d, J = 6.8 Hz), 1.00 (d, J = 6.8 Hz), 1.04 (d, J = 6.8 Hz), 6H]; [1.52 - 1.64 (m); 1.72 - 1.85 (m); 1H]; [2.05 (s), 2.14 (s), 3H]; 3.80 (s, 3H); 4.00 - 4.15 (m, 1H); [4.17 (s), 4.22 (s),
25 2H]; 4.95 - 5.15 (m, 2H); 5.04 (d, J = 11.2 Hz, 0.5H); 5.47 (d, J = 10.7 Hz, 0.5H); [6.10 (d, J = 5.5 Hz); 6.12 (d, J = 7.3 Hz), 1H]; 6.80 - 6.93 (m, 3H); 7.22 - 7.41 (m, 6H).

30 d. 1-[3-[5-(3-Methoxybenzyl)]-1,2,4-oxadiazolyl]-2-(S)-Benzyloxycarbonylamino-3-Methyl-Butan-1-ol.

 This compound was prepared using the hydrolysis conditions in Example I but with the intermediate 1-[3-[5-(3-methoxybenzyl)]-1,2,4-oxadiazolyl]-1-acetoxy-
35 2-(S)-benzyloxycarbonylamino-3-methyl-butane, oil (yield 83.6 %).

¹H (CDCl₃) δ [0.93 (d, J = 6.7 Hz); 0.98 (d, J = 6.7 Hz); 1.00 (d, J = 6.7 Hz); 1.08 (d, J = 6.7 Hz);

6H]; [1.59 - 1.72 (m); 1.92 - 2.06 (m); 1H]; [3.02 (d, J = 8.1 Hz), 3.21 (d, J = 5.7 Hz), 1H]; [3.79 (s), 3.80 (s), 3H]; [3.76 - 3.86 (m), 3.92 - 4.02 (m), 1H]; [4.17 (s), 4.20 (s); 2H]; 4.98 - 5.10 (m, 1.5H); [5.01 (ABq, J = 12.5 Hz), 5.11 (ABq, J = 12.1 Hz), 2H]; 5.28 (d, J = 10.4 Hz, 0.5H); 6.80 - 6.91 (m, 3 H); 7.22 - 7.40 (m, 6H).

e. 1-[3-[5(3-methoxybenzyl)]-1,2,4-oxadiazolyl]-2-(S)-Amino-3-Methyl-Butan-1-ol Hydrochloride.

This compound was prepared following substantially the same procedure as described in Example I utilizing 1-[3-[5-(3-methoxybenzyl)]-1,2,4-oxadiazolyl]-2-(S)-benzyloxycarbonylamino-3-methylbutan-1-ol; heavy oil (yield 90.7%).

^1H (CDCl_3) δ 0.80 - 1.12 (m, 6H); [1.84 - 1.96 (m); 1.96 - 2.10 (m), 1H]; 3.60 - 3.75 (m, 1H); [3.74 (s), 3.76 (s), 3H]; 4.05 - 4.30 (m, 2H); [5.26 (d, J = 8.4 Hz), 5.35 (d, J = 3.1 Hz), 1H]; 6.76 - 6.90 (m, 3H); 7.16 - 7.22 (m, 1H); 8.08 - 8.25 (2brs, 3H).

f. (Benzyloxycarbonyl)-L-Valyl-N-[1-[(3-[5-(3-methoxybenzyl)-1,2,4-oxadiazolyl]hydroxymethyl)-2-(S)-Methylpropyl]-L-Prolinamide.

This compound was prepared by a similar coupling procedure as described in Example I using the hydrochloride of 1-[3-[5-(3-methoxybenzyl)]-1,2,4-oxadiazolyl]-2-(S)-amino-3-methyl-butan-1-ol to give (benzyloxycarbonyl)-L-valyl-N-[1-[(3-[5-(3-phenylethyl)-1,2,4-oxadiazolyl]hydroxymethyl)-2-(S)-methylpropyl]-L-prolinamide as a white solid (yield 50.8 %).

^1H (CDCl_3) δ 0.76 - 1.12 (m, 12H); 1.65 - 2.25 (m, 6H); 3.45 - 3.80 (m, 2H); 3.77 (s, 3H); 3.85 - 4.05 (m, 0.5H); [4.10 (s), 4.18 (s), 2H]; 4.15 - 4.65 (m, 3H); 4.75 - 5.15 (m, 2.5H); [5.70 (d, J = 9.1 Hz), 5.71 (d, J = 8.9 Hz), 0.5H]; 6.11 (d, J = 9.1 Hz, 0.5H); 6.75 - 7.00 (m, 3.5H), 7.15 - 7.45 (m, 7.5H).

EXAMPLE IX

(Benzyloxycarbonyl)-L-Valyl-N-[1-[(3-[5-(2,6-difluorobenzyl)-1,2,4-oxadiazolyl]carbonyl]-2-(S)-Methylpropyl]-L-Prolinamide.

5 The compound was prepared using a similar oxidative procedure as described in Example I but utilizing (benzyloxycarbonyl)-L-valyl-N-[1-[(3-[5-(2,6-difluorobenzyl)-1,2,4-oxadiazolyl]hydroxymethyl)-2-(S)-methylpropyl]-L-prolinamide for the alcohol; R_f = 0.24 (silica gel 1:1, ethyl acetate:hexane). The material was purified by column chromatography on silica gel (1:1, ethyl acetate:hexane) to give a white solid, 49.2 %yield.

^1H NMR (CDCl_3) δ [0.87 (d, J = 6.8 Hz); 0.94 (d, J = 6.8 Hz); 1.02 (d, J = 6.8 Hz), 12H]; 1.80 - 2.20 (m, 4H); 2.20-2.45 (m, 2H); 3.55-3.65 (m, 1H); 3.65 - 3.80 (m, 1H); 4.25 - 4.45 (m, 1H); 4.37 (s, 2H), 4.61 (dd, J_1 = 2.5 Hz; J_2 = 7.8 Hz; 1H); 5.09 (ABq, J = 12.2 Hz, 2H); 5.28 (dd, J_1 = 4.9 Hz, J_2 = 7.3 Hz, 1H); 5.53 (d, J = 9.3 Hz, 1H); 6.95 (t, J = 8.5 Hz, 1H); 7.29 - 7.40 (m, 8H).

^{13}C NMR (CDCl_3) δ 17.85, 17.56, 19.45, 19.79, 20.39, 25.11, 27.18, 30.31, 31.42, 47.74, 57.47, 59.71, 61.46, 66.92, 111.50 (dd, J_1 = 7.0 Hz, J_2 = 25.0 Hz), 128.00, 128.10, 128.48, 130.11 (t, J = 10.3 Hz), 136.28, 156.41, 161.30 (dd, J_1 = 7.2 Hz, J_2 = 249.8 Hz), 164.86, 171.07, 172.16, 177.90, 189.90.

IR (Deposit) 3430, 3028, 1717, 1684, 1629, 1577, 1509, 1436 cm^{-1} .

$\text{C}_{32}\text{H}_{37}\text{N}_5\text{O}_6 \cdot 0.5 \text{ H}_2\text{O}$

	%C	%H	%N
Theory	60.56	6.03	11.03
Found	60.53	6.15	10.90

The intermediate (benzyloxycarbonyl)-L-valyl-N-[1-[(3-[5-(2,6-difluorobenzyl)-1,2,4-

oxadiazolyl]hydroxymethyl]-2-(S)-methylpropyl]-L-prolinamide was prepared as follows:

- a. 1-[3-[5-(2,6-Difluorobenzyl)]-1,2,4-oxadiazolyl]-
5 1-Acetoxy-2-(S)-Benzyloxycarbonylamino-3-Methyl-Butane.

This compound was prepared using a similar cyclization procedure as reported in Example IV except 2,6-difluorophenylacetic anhydride was used, yellow
10 oil; TLC R_f = 0.58, silica gel, 1:1 (ethyl acetate:hexane).

^1H (CDCl_3) δ 0.90 - 1.01 (m, 6H); 1.50 - 2.00 (m, 1H), [2.05 (s), 2.10 (s), 3H]; [3.76 (s), 3.86 (s), 2H]; 3.95 - 4.15 (m, 1H); 4.95 - 5.20 (m, 2.5H); 5.40
15 - 5.50 (brd, 0.5H); 6.00 - 6.15 (brs, 1H); 6.86 - 6.93 (m, 3H); 7.25 - 7.36 (m, 5H).

- b. 1-[3-[5-(2,6-Difluorobenzyl)]-1,2,4-oxadiazolyl]-
2-(S)-Benzyloxy carbonylamino-3-Methyl-Butan-1-ol.

20 This compound was prepared using the hydrolysis conditions in Example I but with the intermediate 1-[3-[5-(2,6-difluorobenzyl)]-1,2,4-oxadiazolyl]-1-acetoxy-2-(S)-benzyloxycarbonylamino-3-methyl-butane, oil (yield 49.8 %); TLC R_f = 0.47, silica gel, 1:1
25 (ethyl acetate:hexane).

^1H (CDCl_3) δ [0.92 (d, J = 6.6 Hz), 0.95 (d, J = 6.6 Hz), 0.96 (d, J = 6.6 Hz), 1.04 (d, J = 6.6 Hz), 6H]; 1.50 - 1.70 (m, 0.5H); 1.80 - 2.05 (m, 0.5H); [3.07 (d, J = 8.1 Hz), 3.32 (d, J = 6.2 Hz), 1H];
30 [3.70 - 3.82 (m), 3.90 - 3.98 (m), 1H]; [4.25 (s), 4.28 (s); 2H]; 4.98 - 5.15 (m, 3.5H); 5.27 (d, J = 9.9 Hz, 0.5H); 6.89 - 6.95 (m, 2H); 7.20 - 7.38 (m, 6H).

- c. 1-[3-[5-(2,6-Difluorobenzyl)]-1,2,4-oxadiazolyl]-
35 2-(S)-Amino-3-Methyl-Butan-1-ol Hydrochloride.

This compound was prepared following substantially the same procedure as described in Example I utilizing 1-[3-[5-(2,6-difluorobenzyl)]-

1,2,4-oxadiazolyl]-2-(S)-benzyloxycarbonylamino-3-methyl-butan-1-ol; Heavy oil (yield 90.5 %).

¹H (DMSO-d₆) δ 0.90 - 1.02 (m, 6H); 1.70 - 1.95 (m, 1H); 3.10 - 3.35 (m, 1H); 4.38 (s, 2H); 4.80 - 4.95 (m, 0.5H); 5.00 - 5.10 (m, 0.5H); 7.00 - 7.50 (m, 3H); 8.01 (brs, 3H).

d. (Benzyloxycarbonyl)-L-Valyl-N-[1-[(3-[5-(2,6-Difluorobenzyl)-1,2,4-oxadiazolyl]hydroxymethyl)-2-(S)-Methylpropyl]-L-Prolinamide.

This compound was prepared by a similar coupling procedure as described in Example I using the hydrochloride of 1-[3-[5-(2,6-difluorobenzyl)]-1,2,4-oxadiazolyl]-2-(S)-amino-3-methyl-butan-1-ol to give (benzyloxycarbonyl)-L-valyl-N-[1-[(3-[5-(2,6-difluorobenzyl)-1,2,4-oxadiazolyl]hydroxymethyl)-2-(S)-methylpropyl]-L-prolinamide as a white solid (yield 68.4 %); TLC R_f = 0.17, silica gel, 3:2 (ethyl acetate:hexane).

¹H (CDCl₃) δ 0.80 - 1.10 (m, 12H); [1.70 - 2.20 (m, 5.7H), 2.45 - 2.55 (m, 0.3H)]; 3.50 - 3.65 (m, 1H); 3.66 - 3.80 (m, 1H); 3.85 - 3.94 (m, 1H); 4.05 - 4.66 (m, 4H); [4.80 (d, J = 8.3 Hz), 4.88 (d, J = 12.0 Hz), 1H]; 4.98 - 5.13 (m, 2H); [5.62 (d, J = 9.3 Hz), 5.73 (d, J = 9.2 Hz), 1H]; 6.03 (d, J = 9.3 Hz, 0.5H); 6.83 - 6.97 (m, 1H); 7.17 - 7.35 (m, 8H); 7.72 (d, J = 9.9 Hz, 0.5H).

EXAMPLE X

(Benzyloxycarbonyl)-L-Valyl-N-[1-[(3-[5-(trans-styryl)-1,2,4-oxadiazolyl]carbonyl)-2-(S)-Methylpropyl]-L-Prolinamide.

The compound was prepared using a similar oxidative procedure as described in Example I but utilizing (benzyloxycarbonyl)-L-valyl-N-[1-[(3-[5-(trans-styryl)-1,2,4-oxadiazolyl]hydroxymethyl)-2-(S)-methylpropyl]-L-prolinamide; TLC R_f = 0.25, (1:1, ethyl acetate:hexane). The material was purified by

column chromatography on silica gel (1:1, ethyl acetate:hexane) to give a white solid.

¹H NMR (CDCl₃) δ [0.92 (d, J = 6.9 Hz); 0.96 (d, J = 6.9 Hz); 1.02 (d, J = 6.9 Hz), 1.05 (d, J = 6.9 Hz), 12H]; 1.80 - 2.25 (m, 4H); 2.25 - 2.45 (m, 2H); 3.55 - 3.70 (m, 1H); 3.76 (dd, J₁ = 6.0 Hz, J₂ = 8.3 Hz, 1H); 4.36 (dd, J₁ = 6.6 Hz, J₂ = 8.8 Hz, 1H); 4.65 (dd, J₁ = 2.5 Hz, J₂ = 8.0 Hz, 1H); 5.1 (ABq, J = 12.3 Hz, 2H); 5.36 (dd, J₁ = 5.1 Hz, J₂ = 7.2 Hz, 1H); 5.55 (d, J = 9.2 Hz, 1H); 7.04 (d, J = 16.4 Hz, 1H); 7.30 - 7.70 (m, 11H); 7.98 (d, J = 16.4 Hz, 1H).

¹³C NMR (CDCl₃) δ 17.30, 17.57, 19.51, 19.85, 25.15, 27.19, 30.48, 31.46, 47.78, 57.48, 59.78, 61.53, 66.96, 109.11, 128.02, 128.15, 128.51, 129.14, 131.03, 133.97, 136.29, 143.00, 144.65, 156.43, 165.08, 171.04, 172.22, 176.73, 190.35.

IR (Deposit) 3429, 3028, 1717, 1682, 1641, 1580, 1509, 1437 cm⁻¹.

20 C₃₃H₃₉N₅O₆ · H₂O

	%C	%H	%N
Theory	63.96	6.67	11.30
Found	64.40	6.41	11.29

25 The intermediate (benzyloxycarbonyl)-L-valyl-N-[1-[(3-[5-(trans-styryl)-1,2,4-oxadiazolyl]hydroxymethyl]-2-(S)-methylpropyl]-L-prolinamide was prepared as follows:

30 a. 1-[(N-Hydroxy)carboximid-N-(trans-Cinnamyl)amido]-1-Acetoxy-2-(S)-Benzyloxycarbonylamino-3-Methyl-Butane.

This compound was prepared following substantially the same procedure as described in Example I, except trans-cinnamyl chloride was used as the acid chloride; 48.1%; TLC R_f = 0.42 and 0.31(diastereomers), 3:2 ethyl acetate:hexane.

¹H (CDCl₃) δ [0.93 (d, J = 6.7 Hz), 0.98 (d, J = 6.7 Hz), 1.00 (d, J = 6.7 Hz), 1.04 (d, J = 6.7 Hz), 6H]; [1.95 (s), 2.10 (s), 3H]; 1.90 - 2.02 (m, 1H); 3.90 - 4.04 (m, 1H); 4.89 - 5.31 (m, 5 H); [5.47 (d, J = 7.0 Hz), 5.48 (d, J = 6.8 Hz), 1H]; [6.52 (d, J = 16.0 Hz), 6.56 (d, J = 16.0 Hz), 1H]; 7.28 - 7.55 (m, 10H); [7.79 (d, J = 16.0 Hz), 7.80 (d, J = 16.0 Hz), 1H].

10 b. 1-[3-[5-(trans-Styryl)]-1,2,4-oxadiazolyl]-1-Acetoxy-2-(S)-Benzyloxycarbonylamino-3-Methyl-Butane.

This compound was prepared using a similar cyclization procedure as reported in Example I except 1-[(N-hydroxy)carboximid-N-(trans-cinnamyl)amido]-1-acetoxy-2-(S)-benzyloxycarbonylamino-3-methyl-butane
15 was used; viscous oil (yield 77.3 %), TLC R_f = 0.74 and 0.68 (diastereomers), silica gel, 3:2 (ethyl acetate:hexane).

¹H (CDCl₃) δ [0.96 (d, J = 7.0 Hz), 1.01 (d, J = 7.0 Hz), 1.03 (d, J = 7.0 Hz), 1.05 (d, J = 7.0 Hz), 6H]; [1.54 - 1.66 (m), 1.74 - 1.88 (m), 1H]; [2.06 (s), 2.14 (s), 3H]; 4.04 - 4.20 (m, 1H); 4.98 - 5.20 (m, 2.5H); 5.57 (d, J = 10.5 Hz, 0.5 Hz); [6.11 (d, J = 3.6 Hz), 6.16 (d, J = 4.9 Hz), 1H]; [6.96 (d, J = 16.4 Hz), 7.00 (d, J = 16.4 Hz), 1 H]; 7.28 - 7.61 (m, 10H); [7.80 (d, J = 16.4 Hz), 7.84 (d, J = 16.4 Hz), 1H].

c. 1-[3-[5-(trans-Styryl)]-1,2,4-oxadiazolyl]-2-(S)-Benzyloxy carbonylamino-3-Methyl-Butan-1-ol.

This compound was prepared using the hydrolysis conditions in Example I but with the intermediate 1-[3-[5-(trans-styryl)]-1,2,4-oxadiazolyl]-1-acetoxy-2-(S)-benzyloxycarbonylamino-3-methyl-butane, oil (yield
35 92.0 %); TLC R_f = 0.61, silica gel, 1:1 (ethyl acetate:hexane).

¹H (CDCl₃) δ [0.96 (d, J = 6.7 Hz), 1.01 (d, J = 6.7 Hz), 1.09 (d, J = 6.7 Hz), 6H]; [1.60 - 1.80 (m),

1.95 - 2.15 (m), 1H]; [3.23 (d, J = 8.1 Hz), 3.47 (d, J = 6.0 Hz), 1H]; [3.80 - 3.95 (m), 3.95 - 4.10 (m), 1H]; 5.00 - 5.42 (m, 3.5H); 5.40 (d, J = 10.3 Hz, 0.5 Hz); [6.95 (d, J = 16.4 Hz), 6.98 (d, J = 16.4 Hz), 1H]; 7.28 - 7.62 (m, 10H); [7.80 (d, J = 16.4 Hz), 7.82 (d, J = 16.4 Hz), 1H].

d. 1-[3-[5-(trans-Styryl)]-1,2,4-oxadiazolyl]-2-(S)-Amino-3-Methyl-Butan-1-ol Hydrochloride.

10 This compound was prepared following substantially the same procedure as described in Example I utilizing 1-[3-[5-(trans-styryl)]-1,2,4-oxadiazolyl]-2-(S)-benzyloxycarbonylamino-3-methylbutan-1-ol; yellow solid (yield 73.0 %).

15 ^1H (CDCl₃) δ [1.04 (d, J = 6.7 Hz), 1.13 (d, J = 6.7 Hz), 1.15 (d, J = 6.7 Hz), 1.19 (d, J = 6.7 Hz), 6H]; 1.85 - 2.20 (m, 1H); 3.55 - 3.70 (m, 1H); [5.26 (d, J = 7.2 Hz), 5.41 (d, J = 3.6 Hz), 1H]; 6.45 (brs, 1H); [7.00 (d, J = 16.4 Hz), 7.03 (d, J = 16.4 Hz), 1H]; 7.20 - 7.70 (m, 5H); [7.83 (d, J = 16.4 Hz), 7.91 (d, J = 16.4 Hz), 1H], 8.43 (bs, 3H).

e. (Benzyloxycarbonyl)-L-Valyl-N-[1-[(3-[5-(trans-Styryl)]-1,2,4-oxadiazolyl]hydroxymethyl)-2-(S)-Methylpropyl]-L-Prolinamide.

25 This compound was prepared by a similar coupling procedure as described in Example I using the hydrochloride of 1-[3-[5-(trans-styryl)]-1,2,4-oxadiazolyl]-2-(S)-amino-3-methylbutan-1-ol to give (benzyloxycarbonyl)-L-valyl-N-[1-[(3-[5-(trans-styryl)]-1,2,4-oxadiazolyl]hydroxymethyl)-2-(S)-methylpropyl]-L-prolinamide as a white solid (yield 66.5%); TLC R_f = 0.34 and 0.27 (diastereomers), silica gel, 7:3 (ethyl acetate:hexane).

35 ^1H (CDCl₃) δ 0.87 - 1.10 (m, 12H); 1.65 - 2.65 (m, 6H); 3.50 - 3.85 (m, 2H); 3.90 - 4.95 (m, 4H); 5.00 - 5.20 (m, 2H); [5.39 (d, J = 8.9 Hz), 5.42 (d,

$J = 8.9 \text{ Hz}$), 5.60 (d, $J = 9.2 \text{ Hz}$), 5.82 (d, $J = 9.2 \text{ Hz}$), 1H]; 6.91 - 7.81 (m, 14H).

EXAMPLE XI

- 5 (Benzyloxycarbonyl)-L-Valyl-N-[1-[(3-[5-(*trans*-4-Trifluoro methylstyryl)-1,2,4-oxadiazolyl]carbonyl]-2-(S)-Methylpropyl]-L-Prolinamide.

The compound was prepared using a similar oxidative procedure as described in Example I but
 10 utilizing (benzyloxycarbonyl)-L-valyl-N-[1-[(3-[5-(*trans*-4-trifluoromethylstyryl)-1,2,4-oxadiazolyl]hydroxymethyl]-2-(S)-methylpropyl]-L-prolinamide; TLC $R_f = 0.49$, (3:2, ethyl acetate:hexane). The material was purified by column
 15 chromatography on silica gel (3:2, ethyl acetate:hexane) to give a white solid.

^1H NMR (CDCl_3) δ [0.93 (d, $J = 7.1 \text{ Hz}$); 0.96 (d, $J = 7.1 \text{ Hz}$); 1.02 (d, $J = 7.1 \text{ Hz}$), 1.05 (d, $J = 7.0 \text{ Hz}$), 12H]; 1.80 - 2.25 (m, 4H); 2.25 - 2.50 (m, 2H);
 20 3.55 - 3.70 (m, 1H); 3.70 - 3.85 (m, 1H); 4.36 (dd, $J_1 = 7.3 \text{ Hz}$, $J_2 = 9.2 \text{ Hz}$, 1H); 4.66 (dd, $J_1 = 2.2 \text{ Hz}$, $J_2 = 7.7 \text{ Hz}$, 1H); 5.10 (ABq, $J = 12.5 \text{ Hz}$, 2H); 5.34 (dd, $J_1 = 5.5 \text{ Hz}$, $J_2 = 5.5 \text{ Hz}$, 1H); 5.52 (d, $J = 8.9 \text{ Hz}$, 1H); 7.13 (d, $J = 16.4 \text{ Hz}$, 1H); 7.30 - 7.45 (m,
 25 4H); 7.48 (d, $J = 7.1 \text{ Hz}$, 1H); 7.71 (s, 5H); 7.98 (d, $J = 16.4 \text{ Hz}$, 1H).

^{13}C NMR (CDCl_3) δ 17.35, 17.59, 19.49, 19.83, 25.15, 27.12, 30.42, 31.45, 47.79, 57.50, 59.73, 61.57, 66.96, 111.58, 126.10, 126.15, 128.02, 128.14,
 30 128.27, 128.52, 132.63, 136.28, 137.20, 142.67, 156.43, 165.14, 171.10, 172.29, 176.08, 190.19.

IR (Deposit) 3430, 3026, 1719, 1681, 1641, 1510, 1509 cm^{-1} .

35 $\text{C}_{34}\text{H}_{36}\text{N}_5\text{O}_6 \text{ F}_3$

	%C	%H	%N
Theory	60.98	5.72	10.46
Found	60.88	5.74	10.50

The intermediate (benzyloxycarbonyl)-L-valyl-N-[1-[(3-[5-(trans-4-trifluoromethylstyryl)-1,2,4-oxadiazolyl]hydroxymethyl)-2-methylpropyl]-L-prolinamide was prepared as follows:

5

a. 1-[(N-Hydroxy)carboximid-N-(trans-4-trifluoromethyl cinnamyl)amido]-1-Acetoxy-2-(S)-Benzyloxycarbonylamino-3-Methyl-Butane.

This compound was prepared following

10 substantially the same procedure as described in Example I, except trans-4-trifluoromethylcinnamyl chloride was used as the acid chloride; 78.6 %; TLC R_f = 0.58 and 0.45 (diastereomers), 3:2 ethyl acetate:hexane.

15 ^1H (CDCl_3) δ [0.92 (d, J = 6.7 Hz), 0.97 (d, J = 6.7 Hz), 0.99 (d, J = 6.7 Hz), 1.03 (d, J = 6.7 Hz), 6H]; [1.96 (s), 2.10 (s), 3H]; 1.80 - 2.10 (m, 1H); 3.85 - 4.10 (m, 1H); 4.95 - 5.30 (m, 5H); [5.46 (d, J = 7.1 Hz), 5.47 (d, J = 6.8 Hz), 1H]; [6.60 (d, J =
20 16.0 Hz), 6.64 (d, J = 16.0 Hz), 1H]; 7.24 - 7.42 (m, 4H); 7.55 - 7.70 (m, 5H); [7.78 (d, J = 16.0 Hz), 7.79 (d, J = 16.0 Hz), 1H].

b. 1-[3-[5-(trans-4-Trifluoromethylstyryl)]-1,2,4-oxadiazolyl]-1-Acetoxy-2-(S)-Benzyloxycarbonylamino-3-Methyl-Butane.

This compound was prepared using a similar cyclization procedure as reported in Example I except 1-[(N-hydroxy)carboximid-N-(trans-4-trifluoromethylcinnamyl)amido]-1-acetoxy-2-(S)-benzyloxycarbonylamino-3-methyl-butane was used; yellow solid (yield 72.6 %), TLC R_f = 0.90, silica gel, 1:1 (ethyl acetate:hexane).

30 ^1H (CDCl_3) δ [0.96 (d, J = 6.7 Hz), 1.01 (d, J = 6.7 Hz), 1.05 (d, J = 6.7 Hz), 6H]; [1.54 - 1.66 (m), 1.74 - 1.88 (m), 1H]; [2.07 (s), 2.15 (s), 3H]; 4.04 - 4.20 (m, 1H); [5.03 (s), 5.13 (s), 2H]; [5.07 (d, J = 10.4 Hz), 5.48 (d, J = 10.4 Hz), 1H]; [6.11 (d, J =
35 6.7 Hz), 1.05 (d, J = 6.7 Hz), 6H]; [1.54 - 1.66 (m), 1.74 - 1.88 (m), 1H]; [2.07 (s), 2.15 (s), 3H]; 4.04 - 4.20 (m, 1H); [5.03 (s), 5.13 (s), 2H]; [5.07 (d, J = 10.4 Hz), 5.48 (d, J = 10.4 Hz), 1H]; [6.11 (d, J =

3.5 Hz), 6.16 (d, $J = 5.1$ Hz), 1H]; [7.02 (d, $J = 16.4$ Hz), 7.07 (d, $J = 16.4$ Hz), 1H]; 7.20 - 7.45 (m, 4H); 7.65 - 7.75 (m, 5H); [7.81 (d, $J = 16.4$ Hz), 7.84 (d, $J = 16.4$ Hz), 1H].

5

c. 1-[3-[5-(*trans*-4-Trifluoromethylstyryl)]-1,2,4-oxadiazolyl]-2-(*S*)-Benzyloxycarbonylamino-3-Methyl-Butan-1-ol.

This compound was prepared using the hydrolysis conditions in Example I but with the intermediate 1-[3-[5-(*trans*-4-trifluoromethylstyryl)]-1,2,4-oxadiazolyl]-1-acetoxy-2-(*S*)-benzyloxycarbonylamino-3-methyl-butane, yellow solid (yield 84.1 %); TLC $R_f = 0.62$, silica gel, 1:1 (ethyl acetate:hexane).

^1H (CDCl_3) δ [0.96 (d, $J = 6.6$ Hz), 1.01 (d, $J = 6.6$ Hz), 1.10 (d, $J = 6.6$ Hz), 6H]; [1.60 - 1.80 (m), 1.95 - 2.15 (m), 1H]; [3.42 (d, $J = 8.0$ Hz), 3.61 (d, $J = 6.0$ Hz), 1H]; [3.80 - 3.95 (m), 3.95 - 4.10 (m), 1H]; 5.00 - 5.42 (m, 3H); [5.38 (d, $J = 10.3$ Hz), 5.27 (d, $J = 9.5$ Hz), 1H]; [7.03 (d, $J = 16.4$ Hz), 7.06 (d, $J = 16.4$ Hz), 1H]; 7.20 - 7.45 (m, 4H); 7.65 - 7.75 (m, 5H); [7.80 (d, $J = 16.4$ Hz), 7.83 (d, $J = 16.4$ Hz), 1H].

d. 1-[3-[5-(*trans*-4-Trifluoromethylstyryl)]-1,2,4-oxadiazolyl]-2-(*S*)-Amino-3-Methyl-Butan-1-ol Hydrochloride.

This compound was prepared following substantially the same procedure as described in Example I utilizing 1-[3-[5-(*trans*-4-trifluoromethylstyryl)]-1,2,4-oxadiazolyl]-2-(*S*)-benzyloxycarbonylamino-3-methyl-butan-1-ol; white solid (yield 63.2 %).

^1H (CDCl_3) δ [1.04 (d, $J = 6.6$ Hz), 1.30 - 1.75 (m), 1.17 (d, $J = 6.6$ Hz), 6H]; 1.85 - 2.20 (m, 1H); 3.55 - 3.70 (m, 1H); [5.26 (d, $J = 7.3$ Hz), 5.44 (brd), 1H]; 6.60 (brs, 1H); [7.17 (d, $J = 16.4$ Hz), 7.10 (d, $J = 16.4$ Hz), 1H]; [7.28 (d, $J = 6.5$ Hz),

7.37 (d, $J = 8.3$), 1H]; 7.55 - 7.75 (m, 3H); [7.85 (d, $J = 16.4$ Hz), 7.96 (d, $J = 16.4$ Hz), 1H]; 8.43 (brs, 3H).

- 5 e. (Benzyloxycarbonyl)-L-Valyl-N-[1-[(3-[5-(trans-4-Trifluoro methylstyryl)-1,2,4-oxadiazolyl]hydroxymethyl)-2-(S)-Methylpropyl]-L-Prolinamide.

This compound was prepared by a similar coupling
10 procedure as described in Example I using the hydrochloride of 1-[3-[5-(trans-4-trifluoromethylstyryl)]-1,2,4-oxadiazolyl]-2-(S)-amino-3-methyl-butan-1-ol to give
(benzyloxycarbonyl)-L-valyl-N-[1-[(3-[5-(trans-
15 styryl)-1,2,4-oxadiazolyl]hydroxymethyl)-2-(S)-methylpropyl]-L-prolinamide as a yellow solid (yield 87.5%); TLC $R_f = 0.38$ and 0.29 (diastereomers), silica gel, 7:3 (ethyl acetate:hexane).

^1H (CDCl_3) δ 0.84 - 1.14 (m, 12H); 1.65 - 2.65
20 (m, 6H); 3.50 - 3.85 (m, 2H); 3.902 - 4.90 (m, 4H); 5.00 - 5.20 (m, 2H); [5.56 (d, $J = 9.2$ Hz), 5.88 (d, $J = 9.2$ Hz), 1H]; [6.92 - 7.38 (m), 7.63 - 7.88 (m), 13H).

25 EXAMPLE XII

(Benzyloxycarbonyl)-L-Valyl-N-[1-[(3-[5-(trans-4-Methoxystyryl)-1,2,4-oxadiazolyl]carbonyl)-2-(S)-Methylpropyl]-L-Prolinamide.

The compound was prepared using a similar
30 oxidative procedure as described in Example I but utilizing (benzyloxycarbonyl)-L-valyl-N-[1-[(3-[5-(trans-4-methoxystyryl)-1,2,4-oxadiazolyl]hydroxymethyl)-2-(S)-methylpropyl]-L-prolinamide; TLC $R_f = 0.38$, (3:2, ethyl
35 acetate:hexane). The material was purified by column chromatography on silica gel (3:2, ethyl acetate:hexane) to give a white solid.

¹H NMR (CDCl₃): δ 0.90 (d, J = 6.8 Hz); 0.97 (d, J = 6.8 Hz); 1.03 (d, J = 6.8 Hz), 1.05 (d, J = 6.8 Hz), 12H]; 1.90 - 2.48 (m, 6H); 3.60 - 3.70 (m, 1H); 3.74 - 3.84 (m, 1H); 3.74 - 3.84 (m, 1H); 3.87 (s, 3H); 4.35 (dd, J₁ = 7.0 Hz, J₂ = 9.2 Hz, 1H); 4.62 (dd, J₁ = 2.8 Hz, J₂ = 7.1 Hz, 1H); 5.10 (ABq, J = 12.2 Hz, 2H); 5.39 (dd, J₁ = 4.9 Hz, J₂ = 7.8 Hz, 1H); 5.67 (d, J = 8.9 Hz, 1H); 6.90 (d, J = 16.4 Hz, 1H); 6.96 (d, J = 8.8 Hz, 1H); 7.25 (d, J = 9.0 Hz, 1H); 7.35 (s, 5H); 7.56 (d, J = 8.8 Hz, 2H); 7.93 (d, J = 16.4 Hz, 1H).

¹³C NMR (CDCl₃) δ 17.10, 17.67, 19.42, 19.82, 25.12, 27.60, 30.63, 31.30, 47.94, 55.47, 57.66, 60.11, 61.49, 67.04, 106.44, 114.62, 126.72, 127.96, 128.15, 128.52, 129.97, 144.41, 156.53, 162.04, 164.90, 171.12, 172.53, 177.20, 190.36.

IR (Deposit) 3429, 3031, 2968, 1718, 1682, 1638, 1604, 1512 cm⁻¹.

20 C₃₄H₄₁N₅O₆₇ · 2 H₂O

	%C	%H	%N
Theory	61.16	6.79	10.49
Found	61.59	6.38	10.02

25 The intermediate (benzyloxycarbonyl)-L-valyl-N-[1-[(3-[5-(trans-4-methoxystyryl)-1,2,4-oxadiazolyl]hydroxymethyl)-2-(S)-methylpropyl]-L-prolinamide was prepared as follows:

30 a. 1-[(N-Hydroxy)carboximid-N-(trans-4-methoxycinnamyl)amido]-1-Acetoxy-2-Benzyloxycarbonylamino-3-Methyl-Butane.

This compound was prepared following substantially the same procedure as described in

35 Example I, except trans-4-methoxycinnamyl chloride was used as the acid chloride; 57.5 %; TLC R_f = 0.47 and 0.36 (diastereomers), 3:2 ethyl acetate:hexane.

¹H (CDCl₃) δ [0.92 (d, J = 6.7 Hz), 0.97 (d, J = 6.7 Hz), 0.99 (d, J = 6.7 Hz), 1.04 (d, J = 6.7 Hz), 6H]; [1.95 (s), 2.10 (s), 3H]; 1.88 - 2.25 (m, 1H); 3.84 (s, 3H); 3.90 - 4.10 (m, 1H); 4.90 - 5.33 (m, 5H); [5.47 (d, J = 7.1 Hz), 5.49 (d, J = 6.8 Hz), 1H]; [6.39 (d, J = 15.9 Hz), 6.43 (d, J = 15.9 Hz), 1H]; 6.91 (d, J = 8.7 Hz, 2H); 7.25 - 7.40 (m, 4H); 7.49 (d, J = 8.7 Hz, 1H); 7.50 (d, J = 8.7 Hz, 1H); [7.44 (d, J = 15.9 Hz), 7.75 (d, J = 15.9 Hz), 1H].

10

b. 1-[3-[5-(*trans*-4-Methoxystyryl)]-1,2,4-oxadiazolyl]-1-Acetoxy-2-(S)-Benzyloxycarbonylamino-3-Methyl-Butane.

This compound was prepared using a similar cyclization procedure as reported in Example I except 1-[(N-hydroxy)carboximid-N-(*trans*-4-methoxycinnamyl)amido]-1-acetoxy-2-(S)-benzyloxycarbonylamino-3-methyl-butane was used; yellow solid (yield 53.4 %), TLC R_f = 0.74 and 0.68 (diastereomers), silica gel, 1:1 (ethyl acetate:hexane).

¹H (CDCl₃) δ [0.96 (d, J = 6.6 Hz), 1.00 (d, J = 6.6 Hz), 1.03 (d, J = 6.6 Hz), 1.04 (d, J = 6.6 Hz), 6H]; [1.50 - 1.68 (m), 1.74 - 1.90 (m), 1H]; [2.05 (s), 2.14 (s), 3H]; [3.85 (s), 3.86 (s), 3H]; 4.04 - 4.16 (m, 1H); [4.99 - 5.18 (m), 5.63 (d, J = 10.6 Hz), 3H]; [6.11 (d, J = 3.8 Hz), 6.16 (d, J = 4.9 Hz), 1H]; [6.81 (d, J = 16.3 Hz), 6.85 (d, J = 16.3 Hz), 1H]; [6.94 (d, J = 7.5 Hz, 2H); 6.95 (d, J = 7.5 Hz, 2H); 7.24 - 7.40 (m, 4H); 7.48 - 7.56 (m, 2H); [7.74 (d, J = 16.4 Hz), 7.77 (d, J = 16.4 Hz), 1H].

c. 1-[3-[5-(*trans*-4-Methoxystyryl)]-1,2,4-oxadiazolyl]-2-(S)-Benzyloxycarbonylamino-3-Methyl-Butan-1-ol.

This compound was prepared using the hydrolysis conditions in Example I but with the intermediate 1-[3-[5-(*trans*-4-methoxystyryl)]-1,2,4-oxadiazolyl]-1-

acetoxy-2-(S)-benzyloxycarbonylamino-3-methyl-butane, (yield 85.2 %); TLC R_f = 0.61, silica gel, 3:2 (ethyl acetate:hexane).

^1H (CDCl_3) δ [0.95 (d, J = 6.6 Hz), 1.00 (d, J = 6.6 Hz), 1.09 (d, J = 6.6 Hz), 6 H]; [1.55 - 1.75 (m), 1.95 - 2.15 (m), 1H]; [3.21 (d, J = 8.1 Hz), 3.47 (d, J = 5.4 Hz), 1H]; 3.86 (s, 3H); 3.80 - 4.10 (m, 1H); 5.00 - 5.20 (m, 3H); [5.22 (d, J = 9.8 Hz), 5.43 (d, J = 10.0 Hz), 1H] [6.80 (d, J = 16.4 Hz), 6.83 (d, J = 16.4 Hz), 1H]; 6.91 - 6.99 (m, 2H), [7.22 - 7.44 (m), 7.46 - 7.60 (m), 7H]; [7.74 (d, J = 16.4 Hz), 7.76 (d, J = 16.4 Hz), 1H].

d. 1-[3-[5-(trans-4-Methoxystyryl)]-1,2,4-oxadiazolyl]-2-(S)-Amino-3-Methyl-Butan-1-ol Hydrochloride.

This compound was prepared following substantially the same procedure as described in Example I utilizing 1-[3-[5-(trans-4-methoxystyryl)]-1,2,4-oxadiazolyl]-2-(S)-benzyloxycarbonylamino-3-methyl-butan-1-ol; (yield 93.2 %).

^1H (CDCl_3) δ [1.03 (d, J = 6.9 Hz), 1.12 (d, J = 6.9 Hz), 1.14 (d, J = 6.6 Hz), 6H]; 1.88 - 2.12 (m, 1H); 3.56 - 3.68 (m, 1H); [3.83 (s), 3.84 (s), 3H]; [5.24 (d, J = 7.1 Hz), 5.40 (d, J = 3.7 Hz), 1H]; [6.83 (d, J = 16.2 Hz), 6.87 (d, J = 17.0 Hz), 1H]; [6.91 (d, J = 6.8 Hz), 6.92 (d, J = 7.0 Hz), 2H]; [7.51 (d, J = 8.3 Hz), 7.53 (d, J = 7.0 Hz), 2H]; [7.75 (d, J = 16.3 Hz), 7.83 (d, J = 16.4 Hz), 1H]; 8.42 (brs, 3H).

e. (Benzyloxycarbonyl)-L-Valyl-N-[1-[(3-[5-(trans-4-Methoxystyryl)]-1,2,4-oxadiazolyl]hydroxymethyl)]-2-(S)-Methylpropyl]-L-Prolinamide.

This compound was prepared by a similar coupling procedure as described in Example I using the hydrochloride of 1-[3-[5-(trans-4-methoxystyryl)]-1,2,4-oxadiazolyl]-2-(S)-amino-3-methyl-butan-1-ol to

give (benzyloxycarbonyl)-L-valyl-N-[1-[(3-[5-(trans-methoxystyryl)-1,2,4-oxadiazolyl]hydroxymethyl)-2-methylpropyl]-L-prolinamide as a yellow solid (yield 87.5%); TLC R_f = 0.25, silica gel, 7:3 (ethyl acetate:hexane).

^1H (CDCl_3) δ 0.84 - 1.14 (m, 12H); [1.60 - 2.22 (m), 2.48 - 2.60 (m), 6H]; 3.46 - 3.64 (m, 1H); 3.66 (m, 1H); [3.86 (s), 3.87 (s), 3H]; 3.92 - 4.90 (m, 4H); 5.02 - 5.18 (m, 2H); [5.36 - 5.64 (m), 5.95 (d, J = 9.5 Hz), 1H]; [6.77 (d, J = 16.4 Hz), 6.80 (d, J = 16.4 Hz), 1H]; 6.88 - 7.10 (m, 2H), 7.12 - 7.60 (m, 9H); [7.72 (d, J = 16.4 Hz), 7.73 (d, J = 16.4 Hz), 1H].

15 EXAMPLE XIII

(Benzyloxycarbonyl)-L-Valyl-N-[1-[(3-[5-(3-Thienylmethyl)-1,2,4-oxadiazolyl]carbonyl)-2-(S)-Methylpropyl]-L-Prolinamide.

The compound was prepared using a similar oxidative procedure as described in Example I but utilizing (benzyloxycarbonyl)-L-valyl-N-[1-[(3-[5-(3-thienylmethyl)-1,2,4-oxadiazolyl]hydroxymethyl)-2-(S)-methylpropyl]-L-prolinamide for the alcohol. The material was purified by column chromatography on silica gel (7:3, ethyl acetate:hexane); the material was further purified via RP-HPLC isocratic $\text{CH}_3\text{CN}:\text{H}_2\text{O}$ (60:40), to give the title compound as a white solid after lyophilization; TLC: R_f = 0.51, 7:3, ethyl acetate: hexane.

^1H NMR (CDCl_3) δ [0.88 (d, J = 6.9 Hz); 0.95 (d, J = 6.9 Hz); 1.01 (d, J = 6.9 Hz), 1.02 (d, J = 6.9 Hz), 12H]; 1.84 - 2.20 (m, 4H); 2.20-2.40 (m, 2H); 3.56-3.70 (m, 1H); 3.72 - 3.82 (m, 1H); 4.35 (s, 2H), 4.28 - 4.42 (m, 1H), 4.62 (dd, J_1 = 2.8 Hz; J_2 = 7.8 Hz; 1H); 5.10 (ABq, J = 12.4 Hz, 2H); 5.31 (dd, J_1 = 4.8 Hz, J_2 = 7.3 Hz, 1H); 5.61 (d, J = 8.9 Hz, 1H); 7.08 (d, J = 5.0 Hz, 1H); 7.25 (s, 1H); 7.30 - 7.44 (m, 7H).

^{13}C NMR (CDCl_3) δ 17.12, 17.61, 19.39, 19.78, 25.07, 27.29, 27.67, 30.31, 31.34, 47.79, 57.53, 59.77, 61.44, 66.93, 123.71, 126.70, 127.85, 127.96, 128.09, 128.47, 131.78, 136.22, 156.44, 164.75, 171.13, 172.28, 179.08, 189.95.

IR (Deposit) 3429, 3026, 2968, 1718, 1683, 1635, 1577, 1437 cm^{-1} .

$\text{C}_{32}\text{H}_{39}\text{N}_5\text{O}_6 \cdot 1 \text{ H}_2\text{O}$

	%C	%H	%N
Theory	58.71	6.41	11.41
Found	59.06	6.03	11.31

The intermediate (benzyloxycarbonyl)-L-valyl-N-
[1-[(3-[5-(3-thienylmethyl)-1,2,4-oxadiazolyl]hydroxymethyl)-2-(S)-methylpropyl]-L-prolinamide was prepared as follows:

a. 1-[(N-Hydroxy)carboximid-N-(3-thienylacetyl)amido]-1-Acetoxy-2-(S)-Benzyloxycarbonylamino-3-Methyl-Butane.

This compound was prepared by a similar method as described in Example I but utilizing 3-thienylacetyl chloride as the acid chloride; pale yellow solid (yield 81.0 %); TLC R_f = 0.49 and 0.38 (diastereomers), silica gel, 3:2 (ethyl acetate:hexane).

^1H (CDCl_3) δ [0.89 (d, J = 6.8 Hz), 0.94 (d, J = 6.8 Hz), 0.97 (d, J = 6.8 Hz), 1.00 (d, J = 6.8 Hz), 1.07 (d, J = 6.8 Hz) 6H]; [1.80 - 2.00 (m), 2.04 - 2.18 (m), 1H]; [1.92 (s), 2.08 (s), 3H]; [3.76 (s), 3.80 (s), 2H]; 3.84 - 3.85 (m, 1H); 4.80 - 5.26 (m, 5H); 5.40 (t, J = 7.0 Hz, 1H); [7.02 - 7.10 (m), 7.14 - 7.20 (m), 1H]; 7.28 - 7.40 (m, 8H).

b. 1-[3-[5-(3-Thienylmethyl)]-1,2,4-oxadiazolyl]-1-Acetoxy-2-(S)-Benzyloxycarbonylamino-3-Methyl-Butane.

This compound was prepared using a similar cyclization procedure as reported in Example I except 1-[(N-hydroxy)carboximid-N-(3-thienylacetyl)amido]-1-acetoxy-2-(S)-benzyloxycarbonylamino-3-methyl-butane was used, brown oil (yield 87.3 %) TLC R_f = 0.75, silica gel, 3:2 (ethyl acetate:hexane).

^1H (CDCl_3) δ [0.92 (d, J = 6.7 Hz), 0.99 (d, J = 6.7 Hz), 1.02 (d, J = 6.7 Hz), 6H]; [1.52 - 1.70 (m), 1.72 - 1.81 (m), 1H]; [2.04 (s), 2.13 (s), 3H]; 4.00 - 4.18 (m, 1H); [4.21 (s), 4.26 (s), 2H]; [4.90 - 5.18 (m), 5.44 (d, J = 10.4 Hz, 3H); [6.08 (d, J = 2.4 Hz, 6.10 (d, J = 5.0 Hz, 1H); [7.02 - 7.08 (m), 7.15 - 7.24 (m), 1H]; 7.26 - 7.40 (m, 7H).

c. 1-[3-[5-(3-Thienylmethyl)]-1,2,4-oxadiazolyl]-2-(S)-Benzyloxycarbonylamino-3-Methyl-Butan-1-ol.

This compound was prepared using the hydrolysis conditions in Example I but with the intermediate 1-[3-[5-(3-thienylmethyl)]-1,2,4-oxadiazolyl]-1-acetoxy-2-(S)-benzyloxycarbonylamino-3-methyl-butane, oil (yield 84.7 %); TLC R_f = 0.51, silica gel, 3:2 (ethyl acetate:hexane).

^1H (CDCl_3) δ [0.93 (d, J = 6.7 Hz), 0.97 (d, J = 6.7 Hz), 0.99 (d, J = 6.7 Hz), 1.07 (d, J = 6.7 Hz), 6H]; [1.62 - 1.80 (m); 1.90 - 2.04 (m); 1H]; [3.18 (d, J = 8.1 Hz), 3.37 (d, J = 6.1 Hz), 1H]; 3.74 - 3.84 (m), 3.90 - 4.02 (m), 1H]; [4.20 (s), 4.24 (s); 2H]; [4.94 - 5.19 (m), 5.27 (d, J = 10.4 Hz), 4H] [7.03 (t, J = 4.4 Hz), 7.15 - 7.22 (m), 1H]; 7.27 - 7.42 (m, 7H).

d. 1-[3-[5-(3-Thienylmethyl)]-1,2,4-oxadiazolyl]-2-(S)-Amino-3-Methyl-Butan-1-ol Hydrochloride.

This compound was prepared following substantially the same procedure as described in Example I utilizing 1-[3-[5-(3-thienylmethyl)]-1,2,4-oxadiazolyl]-2-(S)-benzyloxycarbonylamino-3-methyl-butan-1-ol; Heavy oil (yield 95.8%).

¹H (CDCl₃) δ [1.02 (d, J = 6.7 Hz); 1.05 (d, J = 6.7 Hz); 6H]; 1.84 - 2.02 (m, 1H); 3.60 - 3.80 (m, 1H); [4.22 (s), 4.27 (s), 2 H]; [5.25 (d, J = 8.0 Hz), 5.36 (d, J = 3.6 Hz), 1H]; [7.00 (d, J = 5.0 Hz), 7.03 (d, J = 5.0 Hz), 1H]; 7.12 - 7.40 (m, 3H), 8.07 (brs, 3H).

f. (Benzyloxycarbonyl)-L-Valyl-N-[1-[(3-[5-(3-Thienylmethyl)-1, 2,4-oxadiazolyl]hydroxymethyl)-2-(S)-Methylpropyl]-L-Prolinamide.

This compound was prepared by a similar coupling procedure as described in Example I using the hydrochloride of 1-[3-[5-(3-thienylmethyl)-1,2,4-oxadiazolyl]-2-(S)-amino-3-methyl-butan-1-ol to give (benzyloxycarbonyl)-L-valyl-N-[1-[(3-[5-(3-thienylmethyl)-1,2,4-oxadiazolyl]hydroxymethyl)-2-(S)-methylpropyl]-L-prolinamide as a solid (yield 54.5%); TLC R_f = 0.06, silica gel, 7:3 (ethyl acetate:hexane).

¹H (CDCl₃) δ 0.80 - 1.15 (m, 12H); [1.70 - 2.20 (m), 2.44 - 2.56 (m), 6H]; 3.50 - 3.60 (m, 1H); 3.64 - 3.80 (m, 1H); 3.90 - 4.04 (m, 1H); 4.06 - 4.90 (m, 5H); 5.00 - 5.20 (m, 3H); [5.30 - 5.40 (m), 5.55 - 5.67 (m), 1H]; 6.02 (d, J = 9.3 Hz, 0.5H), 6.84 - 7.08 (m, 2H); 7.15 - 7.40 (m, 7H); 7.68 (brd, 0.5H).

EXAMPLE XIV

(Benzyloxycarbonyl)-L-Valyl-N-[1-[(3-[5-(4-Methoxybenzyl)-1, 2,4-oxadiazolyl]carbonyl)-2-(S)-Methylpropyl]-L-Prolinamide.

The compound was prepared using a similar oxidative procedure as described in Example I but utilizing (benzyloxycarbonyl)-L-valyl-N-[1-[(3-[5-(4-methoxybenzyl)-1,2,4-oxadiazolyl]hydroxymethyl)-2-(S)-methylpropyl]-L-prolinamide for the alcohol. The material was purified initially by column chromatography (silica gel, 3:2 ethyl acetate:hexane), and further purified by RP-HPLC (isocratic,

CH₃CN: H₂O: 60:40) to afford an analytical sample of the product as a white solid; $R_f = 0.36$ (silica gel 3:2, ethyl acetate:hexane).

¹H NMR (CDCl₃) δ [0.88 (d, $J = 6.8$ Hz); 0.95 (d, $J = 6.8$ Hz); 1.01 (d, $J = 6.8$ Hz), 12H]; 1.80 - 2.20 (m, 4H); 2.24-2.40 (m, 2H); 3.55-3.65 (m, 1H); 3.67 - 3.78 (m, 1H); 3.80 (s, 3H), 4.24 (s, 2H), 4.35 (dd, $J_1 = 6.4$ Hz; $J_2 = 9.1$ Hz; 1H); 4.62 (dd, $J_1 = 2.6$ Hz, $J_2 = 7.6$ Hz, 1H); 5.10 (ABq, $J = 12.2$ Hz, 2H); 5.30 (dd, $J_1 = 5.2$ Hz, $J_2 = 7.5$ Hz, 1H); 5.49 (d, $J = 9.2$ Hz, 1H); 6.88 (d, $J = 8.7$ Hz, 2H); 7.26 (d, $J = 8.7$ Hz, 2H); 7.36 (s, 5H); 7.38 (d, $J = 8.8$ Hz, 1H).

¹³C NMR (CDCl₃) δ 17.18, 17.54, 19.51, 19.86, 25.15, 27.12, 30.33, 31.44, 32.18, 47.77, 55.31, 57.46, 59.75, 61.46, 66.97, 111.44, 114.45, 124.51, 128.01, 128.14, 128.52, 130.12, 156.40, 158.91, 159.25, 171.00, 172.24, 179.90, 190.05.

IR (Deposit) 3432, 3036, 2969, 1717, 1687, 1633, 1620, 1514 cm⁻¹.

C₃₃H₄₁N₅O₇

	%C	%H	%N
Theory	63.96	6.67	11.30
Found	63.71	6.79	11.59

The intermediate (benzyloxycarbonyl)-L-valyl-N-[1-[(3-[5-(4-methoxybenzyl)-1,2,4-oxadiazolyl]hydroxymethyl)-2-(S)-methylpropyl]-L-prolinamide was prepared as follows:

a. 4-Methoxyphenylacetyl Chloride.

This acid chloride was prepared in a similar manner as for the acid chloride in Example I and was used without further purification.

b. 1-[(N-Hydroxy)carboximid-N-(4-Methoxybenzyl)amido]-1-Acetoxy-2-(S)-Benzyloxycarbonylamino-3-Methyl-Butane.

This compound was prepared following substantially the same procedure as described in Example I, but 4-methoxyphenylacetyl chloride using as the acid chloride; 39.1 %, off-white solid; TLC R_f = 0.51 and 0.43 (diastereomers), silica gel 7:3 (ethyl acetate:hexane).

^1H NMR (CDCl_3) δ [0.89 (d, J = 6.8 Hz); 0.93 (d, J = 6.8 Hz); 0.99 (d, J = 6.8 Hz); 1.06 (d, J = 6.8 Hz); 6H]; [1.91 (s), 2.07 (s), 3H]; 1.80 - 2.20 (m, 1H); [3.66 (s), 3.70 (s), 2H]; [3.78 (s), 3.79 (s), 3H]; 3.80 - 3.95 (m, 1H); [4.80 - 5.24, 5.34 - 5.44 (m), 5H]; [6.83 (d, J = 6.2 Hz), 6.86 (d, J = 6.1 Hz); 2H]; [7.15 - 7.25 (m), 7.18 - 7.40 (m), 3H].

c. 1-[3-[5-(4-Methoxybenzyl)]-1,2,4-oxadiazolyl]-1-Acetoxy-2-(S)-Benzyloxycarbonylamino-3-Methyl-Butane.

This compound was prepared in a similar manner by a cyclization procedure described in Example I utilizing 1-[(N-hydroxy)carboximid-N-(4-methoxybenzyl)amido]-1-acetoxy-2-(S)-benzyloxycarbonylamino-3-methyl-butane; oil (yield 77.2 %), TLC R_f = 0.76, silica gel 3:2 (ethyl acetate:hexane).

^1H (CDCl_3) δ [0.92 (d, J = 6.7 Hz), 0.98 (d, J = 6.7 Hz), 1.01 (d, J = 6.8 Hz), 6H]; [1.50 - 1.64 (m), 1.71 - 1.84 (m), 1H]; [2.03 (s), 2.18 (s), 3H]; [3.77 (s), 3.78 (s), 3H]; 3.96 - 4.22 (m, 3H); [4.84 - 5.18 (m), 5.47 (d, J = 10.4 Hz), 3H]; [6.08 (d, J = 5.3 Hz), 6.10 (d, J = 7.0 Hz), 1H]; 6.80 - 6.92 (m, 2H); 7.16 - 7.42 (m, 7H).

d. 1-[3-[5-(4-Methoxybenzyl)]-1,2,4-oxadiazolyl]-2-(S)-Benzyloxycarbonylamino-3-Methyl-Butan-1-ol.

This compound was prepared using the hydrolysis conditions in Example I but with the intermediate 1-[3-[5-(4-methoxybenzyl)]-1,2,4-oxadiazolyl]-1-acetoxy-2-(S)-benzyloxycarbonylamino-3-methyl-butane, oil

(yield 90.5 %), TLC R_f = 0.58, silica gel 3:2 (ethyl acetate:hexane).

^1H (CDCl_3) δ [0.92 (d, J = 6.7 Hz); 0.96 (d, J = 6.7 Hz); 0.98 (d, J = 6.7 Hz); 1.06 (d, J = 6.7 Hz); 5 6H]; [1.58 - 1.70 (m); 1.82 - 2.04 (m); 1H]; [3.09 (d, J = 8.1 Hz), 3.29 (d, J = 5.6 Hz), 1H]; [3.77 (s), 3.78 (s), 3H]; 3.90 - 4.02 (m, 1H); [4.12 (s), 4.15 (s); 2H]; [4.90 - 5.18 (m), 5.28 (d, J = 10.0 Hz), 4H]; [6.84 (d, J = 8.9 Hz), 6.86 (d, J = 8.7 Hz), 2H]; 10 [7.20 (d, J = 8.5 Hz), 7.22 (d, J = 8.7 Hz), 2H]; 7.28 - 7.39 (m, 5H).

e. 1-[3-[5-(4-Methoxybenzyl)]-1,2,4-oxadiazolyl]-2-(S)-Amino-3-Methyl-Butan-1-ol Hydrochloride.

15 This compound was prepared following substantially the same procedure as described in Example I utilizing 1-[3-[5-(4-methoxybenzyl)]-1,2,4-oxadiazolyl]-2-(S)-benzyloxycarbonylamino-3-methylbutan-1-ol; heavy oil (yield 66.9 %).

20 ^1H (CDCl_3) δ 1.00 (d, J = 6.7 Hz, 3H); 1.04 (d, J = 6.7 Hz, 3H); 1.86 - 2.00 (m, 1H); 3.56 - 3.94 (m, 1H); [3.73 (s), 3.76 (s), 3H]; [4.10 (s), 4.16 (s), 2H]; [5.22 (d, J = 7.8 Hz), 5.31 (d, J = 3.3 Hz), 1H]; [6.81 (d, J = 8.6 Hz), 6.82 (d, J = 8.6), 2H]; [7.15 25 (d, J = 8.6 Hz), 7.19 (d, J = 8.6 Hz), 2H]; 8.03 (brd, 3H).

f. (Benzyloxycarbonyl)-L-Valyl-N-[1-[(3-[5-(4-Methoxybenzyl)]-1, 2,4-oxadiazolyl]hydroxymethyl)-2-(S)-Methylpropyl]-L-Prolinamide .

This compound was prepared by a similar coupling procedure as described in Example I using the hydrochloride of 1-[3-[5-(4-methoxybenzyl)]-1,2,4-oxadiazolyl]-2-(S)-amino-3-methyl-butan-1-ol to give 35 (benzyloxycarbonyl)-L-valyl-N-[1-[(3-[5-(4-methoxybenzyl)]-1,2,4-oxadiazolyl]hydroxymethyl)-2-(S)-methylpropyl]-L-prolinamide as a white solid (yield

50.8 %); TLC R_f = 0.35, silica gel 4:1 (ethyl acetate:hexane).

^1H (CDCl_3) δ 0.77 - 1.12 (m, 12H); [1.70 - 2.20 (m), 2.46 - 2.58 (m), 6H]; 3.46 - 3.74 (m, 2H); [3.76 (s), 3.78 (s), 3H]; 3.88 - 4.90 (m, 6H); 5.00 - 5.20 (m, 2H); [5.53 (d, J = 9.3 Hz), 5.64 (d, J = 9.4 Hz), 1H]; 5.93 (d, J = 9.4 Hz, 0.5H); 6.76 - 6.90 (m, 2H); 7.08 - 7.44 (m, 8H); 7.72 (brd, 0.5H).


10 EXAMPLE XV

(Benzyloxycarbonyl)-L-Valyl-N-[1-[(3-[5-(Phenyl)-1,2,4-oxadiazolyl]carbonyl]-2-(S)-Methylpropyl]-L-Prolinamide.

The compound was prepared using a similar
15 oxidative procedure as described in Example I but utilizing (benzyloxycarbonyl)-L-valyl-N-[1-[(3-[5-(phenyl)-1,2,4-oxadiazolyl]hydroxymethyl]-2-(S)-methylpropyl]-L-prolinamide for the alcohol. Purification was conducted using column chromatography
20 (silica gel, 1:1 to 7:3, ethyl acetate:hexane) followed by purification by RP-HPLC, isocratic $\text{CH}_3\text{CN}:\text{H}_2\text{O}$ (60:40), to give an analytical sample as a white solid after lyophilization; TLC: R_f = 0.42, ethyl acetate: hexane (7:3).

25 ^1H NMR (CDCl_3) δ [0.95 (d, J = 6.8 Hz), 0.99 (d, J = 6.8 Hz), 1.05 (d, J = 6.8 Hz), 1.08 (d, J = 6.8 Hz); 12H]; 1.80 - 2.25 (m, 4H); 2.25-2.50 (m, 2H); 3.55-3.70 (m, 1H); 3.70 - 3.85 (m, 1H); 4.30 - 4.45 (m, 1H), 4.60 - 4.75 (m, 1H); 5.12 (ABq, J = 12.4 Hz,
30 2H); 5.37 - 5.44 (m, 1H), 5.57 (d, J = 9.6 Hz, 1H), 7.25 - 7.70 (m, 10H), 8.62 (d, J = 8.5 Hz, 1H).

^{13}C NMR (CDCl_3) δ 17.32, 17.59, 19.51, 19.86, 25.16, 27.23, 30.49, 31.46, 47.81, 57.51, 59.82, 61.61, 66.98, 123.26, 128.02, 128.14, 129.27, 133.53,
35 136.28, 156.44, 165.33, 171.09, 172.27, 177.17, 190.36.



	%C	%H	%N
Theory	63.68	6.55	11.98
Found	63.51	6.45	11.71

5

The intermediate (benzyloxycarbonyl)-L-valyl-N-[1-[(3-[5-(phenyl)-1,2,4-oxadiazolyl]hydroxymethyl)-2-(S)-methylpropyl]-L-prolinamide was prepared as follows:

10

a. 1-[3-[5-(Phenyl)]-1,2,4-oxadiazolyl]-1-Acetoxy-2-(S)-Benzyloxycarbonylamino-3-Methyl-Butane.

This compound was prepared in a similar manner to that in Example IV except benzoic anhydride was used; yield (75.6 %) of product as a viscous oil.

^1H (CDCl_3) δ [0.97 (d, J = 6.7 Hz), 1.02 (d, J = 6.7 Hz), 1.04 (d, J = 6.7 Hz), 1.06 (d, J = 6.7 Hz) 6H]; [1.56-1.70 (m), 1.76 - 1.90 (m), 1H]; [2.06 (s), 2.15 (s), 3H]; 4.10 - 4.21 (m, 1H); 4.96 - 5.20 (m, 2.5H); 5.61 (d, J = 10.0 Hz, 0.5H); [6.16 (d, J = 3.5 Hz), 6.21 (d, J = 4.9 Hz), 1H]; 7.25 - 7.45 (m, 4H); 7.45 - 7.70 (m, 4H); 8.05 - 8.20 (m, 2H).

b. 1-[3-[5-(Phenyl)]-1,2,4-oxadiazolyl]-2-(S)-Benzyloxycarbonylamino-3-Methyl-Butan-1-ol.

This compound was prepared using hydrolysis conditions in Example I but with the intermediate 1-[3-[5-(phenyl)]-1,2,4-oxadiazolyl]-1-acetoxy-2-(S)-benzyloxycarbonylamino-3-methyl-butane; oil (yield 53.3 %), TLC: R_f = 0.47 and 0.33 (diastereomers), ethyl acetate: hexane (1:1).

^1H (CDCl_3) δ [0.95 (d, J = 7.0 Hz), 0.99 (d, J = 7.0 Hz), 1.01 (d, J = 7.0 Hz), 1.09 (d, J = 7.0 Hz), 6H]; [1.71 - 1.85 (m), 1.94 - 2.09 (m), 1H]; 3.87 - 3.98 (m, 0.5H); 4.03 - 4.16 (m, 0.5H), 4.96 - 5.22 (m, 3.5H), 5.62 (d, J = 9.3 Hz, 0.5H); 7.10 - 7.35 (m, 4H), 7.35 - 7.65 (m, 4H); 7.95 - 8.15 (m, 2H).

c. 1-[3-[5-(Phenyl)]-1,2,4-oxadiazolyl]-2-(S)-Amino-3-Methyl-Butan-1-ol Hydrochloride.

This compound was prepared following substantially the same procedure as described in
5 Example I utilizing 1-[3-[5-(phenyl)]-1,2,4-oxadiazolyl]-2-(S)-benzyloxycarbonylamino-3-methylbutan-1-ol; Heavy oil (yield 54.7 %);

^1H (CDCl_3) δ 1.00-1.35 (m, 6H); 1.85 - 2.15 (m, 1H); 3.75 - 3.90 (m, 0.5H), 3.90 - 4.05 (m, 0.5H);
10 [5.40 (d, J = 8.9 Hz), 5.57 (d, J = 2.6 Hz), 1H]; 7.30 - 7.60 (m, 3H); 7.95 (d, J = 8.0 Hz, 1H); 8.09 (d, J = 7.7 Hz, 1H); 8.17 (brs, 3H).

d. (Benzyloxycarbonyl)-L-Valyl-N-[1-[(3-[5-(Phenyl)]-1,2,4-oxadiazolyl]hydroxymethyl)-2-(S)-Methylpropyl]-L-Prolinamide.

This compound was prepared by a similar coupling procedure as described in Example I using the hydrochloride of 1-[3-[5-(phenyl)]-1,2,4-oxadiazolyl]-
20 2-(S)-amino-3-methyl-butan-1-ol to give (benzyloxycarbonyl)-L-valyl-N-[1-[(3-[5-(phenyl)]-1,2,4-oxadiazolyl]hydroxymethyl)-2-methylpropyl]-L-prolinamide as a white solid (yield 77.2 %); R_f = 0.41, 4:1 ethyl acetate:hexane.

^1H (CDCl_3) δ 0.80 - 1.20 (m, 12H); 1.65 - 2.65 (m, 6H); 3.45 - 3.85 (m, 2H); 3.90 - 4.05 (m, 0.5H); 4.05 - 4.65 (m, 3H); 4.70 - 5.20 (m, 3.5H); [5.21 (t, J = 9.7 Hz), 5.38 (d, J = 9.2 Hz), 1H]; 7.14 (d, J = 9.5 Hz, 0.5H); 7.30 - 7.64 (m, 9.5H), [8.10 (d, J =
30 8.0 Hz), 8.19 (d, J = 8.0 Hz), 1H].

EXAMPLE XVI

(Benzyloxycarbonyl)-L-Valyl-N-[1-[(3-[5-(3-Phenylpropyl)-1,2,4-oxadiazolyl]carbonyl)-2-(S)-
35 Methylpropyl]-L-Prolinamide.


The compound was prepared using a similar oxidative procedure as described in Example I but utilizing (benzyloxycarbonyl)-L-valyl-N-[1-[(3-[5-(3-

phenylpropyl)-1,2,4-oxadiazolyl]hydroxymethyl]-2-(S)-methylpropyl]-L-prolinamide for the alcohol; an analytical sample was obtained from RP-HPLC (isocratic, CH₃CN: H₂O: 60:40) as a white solid; TLC R_f = 0.29, 3:2 (ethyl acetate:hexane).

¹H NMR (CDCl₃) δ [0.90 (d, J = 6.9 Hz); 0.96 (d, J = 6.9 Hz); 1.01 (d, J = 6.9 Hz), 1.03 (d, J = 6.9 Hz), 12H]; 1.82 - 2.40 (m, 6H); 2.20 (p, J = 7.6 Hz, 2H); 2.74 (t, J = 7.6 Hz, 2H); 2.96 (t, 7.6 Hz, 2H); 3.58 - 3.68 (m, 1H); 3.70 - 3.82 (m, 1H); 4.36 (dd, J₁ = 6.8 Hz; J₂ = 9.0 Hz; 1H); 4.60 - 4.63 (m, 1H); 5.10 (ABq, J = 12.3 Hz, 2H); 5.34 (dd, J = 4.9 Hz, J₂ = 9.0 Hz, 1H); 5.75 (d, J = 9.1 Hz, 1H); 7.15 - 7.40 (m, 10H); 7.48 (d, J = 7.4 Hz, 1H)

¹³C NMR (CDCl₃) δ 17.17, 17.59, 19.40, 19.77, 25.07, 27.78, 27.26, 27.76, 30.35, 31.39, 34.81, 47.73, 57.51, 59.71, 61.41, 66.88, 126.26, 127.96, 128.06, 128.41, 128.44, 128.50, 136.29, 140.23, 156.43, 164.71, 171.13, 172.14, 181.27, 190.17.

IR (Deposit) 3297.6, 2966.0, 1718.9, 1680.6, 1632.4 cm⁻¹



	%C	%H	%N
25 Theory	66.11	7.02	11.34
Found	65.87	7.25	11.27

The intermediate (benzyloxycarbonyl)-L-valyl-N-[1-[(3-[5-(3-phenylpropyl)-1,2,4-oxadiazolyl]hydroxymethyl]-2-(S)-methylpropyl]-L-prolinamide was prepared as follows:

a. 4-Phenylbutyryl Chloride.

The acid chloride was prepared in a similar manner as described in Example II except 4-phenylbutyric acid was used, the material was used without further purification.

b. 1-[(N-Hydroxy)carboximid-N-(4-phenylbutyryl)amido]-1-Acetoxy-2-(S)-Benzyloxycarbonylamino-3-Methyl-Butane.

This compound was prepared following

5 substantially the same procedure as described in Example I, except 4-Phenylbutyryl chloride was used as the acid chloride; 34.1%, white solid, TLC R_f = 0.41 and 0.33 (diastereomers), 1:1 (ethyl acetate:hexane).

^1H NMR (CDCl_3) δ [0.91 (d, J = 6.8 Hz); 0.95 (d, J = 6.8 Hz); 0.98 (d, J = 6.8 Hz); 1.02 (d, J = 6.8 Hz); 6H]; [1.94 (s), 2.09 (s), 3H]; 1.84 - 2.12 (m, 3H); 2.36 - 2.47 (m, 2H); 2.64 - 2.73 (m, 2H); 3.84 - 3.96 (m, 1H); 4.76 - 5.24 (m, 5H); 5.30 - 5.45 (m, 1H); 7.14 - 7.40 (m, 10H).

15

c. 1-[3-[5-(3-Phenylpropyl)]-1,2,4-oxadiazolyl]-1-Acetoxy-2-(S)-Benzyloxycarbonylamino-3-Methyl-Butane.

This compound was prepared by a cyclization procedure as described in Example I utilizing 1-[(N-hydroxy)carboximid-N-(3-phenylpropionyl)amido]-1-acetoxy-2-(S)-benzyloxycarbonylamino-3-methyl-butane; oil (yield 85.7%), TLC R_f = 0.63 and 0.60 (diastereomers), 1:1 (ethyl acetate:hexane).

^1H (CDCl_3) δ [0.94 (d, J = 6.7 Hz); 0.99 (d, J = 6.7 Hz); 1.00 (d, J = 6.7 Hz); 1.03 (d, J = 6.7 Hz); 6H]; [1.50 - 1.62 (m); 1.72 - 1.86 (m); 1H]; [2.05 (s), 2.13 (s); 3H]; 2.04 - 2.20 (m, 2H); 2.70 (q, J = 7.7 Hz, 2H); [2.84 (t, J = 7.7 Hz), 2.90 (t, J = 7.7 Hz), 2H]; 4.00 - 4.13 (m, 1H); [5.00 (s), 5.11 (s); 2H]; [5.05 (d, J = 10.4 Hz), 5.50 (d, J = 10.6 Hz); 1H]; [6.08 (d, J = 3.3 Hz); 6.11 (d, J = 5.0 Hz); 1H]; 7.16 - 7.40 (m, 10H).

d. 1-[3-[5-(3-Phenylpropyl)]-1,2,4-oxadiazolyl]-2-(S)-Benzyloxycarbonylamino-3-Methyl-Butan-1-ol.

The compound was prepared using the hydrolysis conditions in Example I with 1-[3-[5-(3-phenylpropyl)]-1,2,4-oxadiazolyl]-1-Acetoxy-2-(S)-

Benzyloxycarbonylamino-3-Methyl-Butane; oil (yield 76.7%), TLC R_f = 0.44, 1:1 (ethyl acetate:hexane).

^1H (CDCl_3) δ [0.94 (d, J = 6.7 Hz); 0.98 (d, J = 6.7 Hz); 0.99 (d, J = 6.7 Hz); 1.08 (d, J = 6.7 Hz); 1.55 - 1.75 (m), 1.90 - 2.15 (m), 1H]; 2.13 (p, J = 7.3 Hz, 2H); 2.69 (q, J = 7.3 Hz, 2H); [2.84 (t, J = 7.3 Hz), 2.87 (t, J = 7.3 Hz), 2H]; [3.22 (d, J = 8.0 Hz), 3.46 (d, J = 6.0 Hz), 1H]; [3.76 - 3.86 (m), 3.92 - 4.02 (m), 1H]; 4.96 - 5.18 (m, 3H); [5.21 (d, J = 9.5 Hz), 5.35 (d, J = 10.0 Hz), 1H]; 7.14 - 7.40 (m, 10H).

e. 1-[3-[5-(3-Phenylpropyl)]-1,2,4-oxadiazolyl]-2-(S)-Amino-3-Methyl-Butan-1-ol Hydrochloride.

This compound was prepared following substantially the same procedure as described in Example I utilizing 1-[3-[5-(3-phenylpropyl)]-1,2,4-oxadiazolyl]-2-(S)-benzyloxycarbonylamino-3-methylbutan-1-ol; heavy oil (yield 29.1 %).

^1H (CDCl_3) δ 1.10 - 1.80 (m, 6H); 1.80 - 2.05 (m, 1H); 2.00 - 2.20 (m, 2H); 2.68 (q, J = 7.3 Hz, 2H); 2.77 - 2.92 (m, 2H); [5.26 (d, J = 8.8 Hz), 5.42 (d, J = 3.0 Hz), 1H]; 7.11 - 7.33 (m, 5H); 8.10 - 8.23 (m, 3H).

f. (Benzyloxycarbonyl)-L-Valyl-N-[1-[(3-[5-(3-Phenylpropyl)-1,2,4-oxadiazolyl]hydroxymethyl)-2-(S)-Methylpropyl]-L-Prolinamide.

This compound was prepared by a similar coupling procedure as described in Example I using the hydrochloride of 1-[3-[5-(3-phenylpropyl)]-1,2,4-oxadiazolyl]-2-(S)-amino-3-methyl-butan-1-ol to give (benzyloxycarbonyl)-L-valyl-N-[1-[(3-[5-(3-phenylpropyl)-1,2,4-oxadiazolyl]hydroxymethyl)-2-methylpropyl]-L-prolinamide as a white foamy solid (yield 60.3 %).

^1H (CDCl_3) δ 0.80 - 1.15 (m, 12H); 1.70 - 2.30 (m, 8H); 2.55 - 2.75 (m, 2H); 2.75 - 3.00 (m, 2H);

3.45 - 3.80 (m, 2H); 3.90 - 4.05 (m, 0.5H), 4.10 - 4.65 (m, 3H), 4.70 - 5.20 (m, 3.5H); [5.51 (t, $J = 8.1$ Hz), 5.82 (d, $J = 9.4$ Hz), 1H]; 6.92 (d, $J = 9.3$ Hz, 0.5H); 7.06 - 7.40 (m, 10.5H).

5

EXAMPLE XVII - Selectivity Studies

The K_i values for different enzymes were determined as described above. The assay buffer was: 0.1 M HEPES, 0.1 M NaCl, 0.01 CaCl_2 , 0.005% Triton X-100, 5% DMSO, pH 7.6. The chromogenic substrates were: Suc-Ala-Ala-Pro-Leu-pNA for α -chymotrypsin (bovine) and porcine pancreatic elastase, and Suc-Ala-Ala-Pro-Phe-pNA for human neutrophil cathepsin G.

15

Enzyme	K_i [μM]	Ratio
HLE ^a	0.0003	1
PPE ^b	0.0149	49.7
α -CH ^b	0.32	1068
Cat-G ^b	>> 30	> 1.0×10^5

20

Buffers: ^a 0.05 M sodium phosphate, 0.1 M NaCl, 0.005% Tx-100, 5% DMSO, pH 7.5; ^b 0.1 M HEPES, 0.1 NaCl, 0.01 M CaCl_2 , 0.005% Tx-100, 5.0% DMSO, pH 7.6.

25

Selectivity of Compound CE-2048: Porcine Pancreatic Elastase, Bovine α -Chymotrypsin and Human Cathepsin G (as compared to Human Leukocyte Elastase)

30 EXAMPLE XVIII - Stability studies (CE-2039, CE-2048 and CE-2049)

Stability of compounds in human plasma was determined as follows. Human plasma was obtained from two male and two female volunteers and previously

stored frozen. Plasma was "spiked" with the compounds to a concentration of 0.025 nM. Samples were incubated 0, 3, and 6 hours at 37°. At each point protein was precipitated by addition of acetonitrile made 0.1 N HCl (3 parts of solution per part of sample). Samples were subjected to centrifugation (15-20 min at 14,000 rpm) and analyzed (200 µL) with reverse-phase HPLC using a 18-90% acetonitrile gradient in 0.1% TFA.


EXAMPLE XIX - K_i Values






Enzyme was added to a mixture of nitroanilide substrate and inhibitor. The rate of aminolysis declined quickly to the steady-state level (V_s). Nitroaniline release from the substrate was monitored at 400-410 nm in an HP-5482 diode-array spectrophotometer. The K_i^{app} values were calculated by a non-linear regression analysis program (ENZFITTER, Elsevier-Biosoft) using the following equations:





$$V_s = V_0 / (1 + [I] / K_i^{app})$$

$$K_i^{app} = K_i (1 + [S] / K_m)$$





where [I] and [S] are the concentrations of inhibitor and substrate, respectively, V_0 and V_s are the velocities of aminolysis in the absence and presence of inhibitor, respectively, K_i^{app} and K_i are apparent and true inhibition constants, respectively, and K_m is the Michaelis constant. The inhibition assays were performed in 0.05 M sodium phosphate, 0.1 M NaCl, 0.005% Triton X-100, 5% DMSO, pH 7.6. The chromogenic substrates were: MeOSuc-Lys (pic) -Ala-Pro-Val-pNA and MeOSuc-Ala-Ala-Pro-Val-pNA.







R ₁	Heterocycle	K _i (nM)
trifluoromethyl		77
methyl		17
difluoromethyl		7.2
2,6-difluorobenzyl		1.7
benzyl		2.0

R_1	Heterocycle	K_1 (nM)
2-thienylmethyl		2.4
styryl		7.9
4-trifluoromethylstyryl		11.5
4-methoxystyryl		8.4

5

R_1	Heterocycle	K_i (nM)
4-methoxybenzyl		1.04
phenyl		9.9
2-phenylethyl		3.8
3-methoxybenzyl		0.5

R_1	Heterocycle	K_i (nM)
3-phenylpropyl		1.84
3-trifluoromethylbenzyl		0.20
2-methoxybenzyl		2.5
2-trifluoromethylphenyl chloromethyl		1.56

5

SEQUENCE LISTING

(1) GENERAL INFORMATION:

- 5 (i) APPLICANT: CORTECH, INC.
- (ii) TITLE OF INVENTION: SUBSTITUTED HETEROCYCLIC
COMPOUNDS USEFUL AS INHIBITORS OF (SERINE
10 PROTEASES) HUMAN NEUTROPHIL ELASTASE
- (iii) NUMBER OF SEQUENCES: 4
- (iv) CORRESPONDENCE ADDRESS:
- 15 (A) NAME: Schwegman, Lundberg & Woessner
(B) STREET: 3500 IDS Center
(C) CITY: Minneapolis
(D) STATE: Minnesota
(E) COUNTRY: USA
(F) POSTAL CODE (ZIP): 55402
20 (G) TELEPHONE: 612-339-0331
(H) TELEFAX: 612-339-3061
- (v) COMPUTER READABLE FORM:
- 25 (A) MEDIUM TYPE: Floppy disk
(B) COMPUTER: IBM PC compatible
(C) OPERATING SYTEM: PC-DOS/MS-DOS
(D) SOFTWARE: PatentIn Release #1.0, Version
#1.30 (EPO)
- 30 (vi) CURRENT APPLICATION DATA:
- (A) APPLICATION NUMBER: UNKNOWN
(B) FILING DATE: 17 NOVEMBER 1995
(C) CLASSIFICATION:
- 35 (vii) ATTORNEY/AGENT INFORMATION:
- (A) NAME: John E. Burke
(B) REGISTRATION NUMBER: 35,836
(C) REFERENCE/DOCKET NUMBER: 392.013WO1
- 40 (2) INFORMATION FOR SEQ ID NO: 1:
- (i) SEQUENCE CHARACTERISTICS:
- (A) LENGTH: 6 amino acids
(B) TYPE: amino acid
45 (C) STRANDEDNESS: single
(D) TOPOLOGY: linear
- (ii) MOLECULE TYPE: peptide
- 50 (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 1:
- Xaa Ala Ala Pro Leu Xaa
1 5

(2) INFORMATION FOR SEQ ID NO: 2:

(i) SEQUENCE CHARACTERISTICS:

- 5 (A) LENGTH: 6 amino acids
(B) TYPE: amino acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: peptide

10

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 2:

Xaa Ala Ala Pro Phe Xaa
1 5

15

(2) INFORMATION FOR SEQ ID NO: 3:

(i) SEQUENCE CHARACTERISTICS:

- 20 (A) LENGTH: 6 amino acids
(B) TYPE: amino acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: peptide

25

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 3:

Xaa Lys Ala Pro Val Xaa
1 5

30

(2) INFORMATION FOR SEQ ID NO: 4:

(i) SEQUENCE CHARACTERISTICS:

- 35 (A) LENGTH: 6 amino acids
(B) TYPE: amino acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: peptide

40

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 4:


Xaa Ala Ala Pro Val Xaa
1 5

45

WHAT IS CLAIMED IS:

1. A compound capable of inhibiting serine proteases according to the formula:

5



10

15

where:


Z is an amino-carbonyl containing group such that the carbon of the heterocycle is attached directly to the carbonyl group of Z;

- 20 R₁ and R₄ are independently selected from alkyl, alkenyl, haloalkyl, haloalkenyl, alkynyl being linear or branched; a phenyl, phenylalkenyl, or phenylalkyl optionally substituted with halogen, cyano, nitro, haloalkyl, amino, aminoalkyl, dialkylamino, alkyl, alkenyl, alkynyl, alkoxy, haloalkoxy, carboxyl, carboalkoxy, alkylcarboxamide, arylcarboxamide, alkylthio, or haloalkylthio groups being linear or branched; a heteroaryl, heteroarylalkyl or
- 25 heteroarylalkenyl wherein the heteroaryl group is a monocyclic five or six membered ring containing one or two heteroatoms selected from oxygen, sulfur and nitrogen in any combination and optionally substituted with halogen, cyano, nitro, haloalkyl, amino, aminoalkyl, dialkylamino, alkyl, alkoxy, alkenyl, alkynyl, haloalkoxy, carboxyl, carboalkoxy,
- 30 alkylcarboxamide, arylcarboxamide, alkylthio or haloalkylthio groups being linear or branched; and

X and Y are O or S.

2. A compound capable of inhibiting serine proteases according to claim 1 wherein:

5 Z is




10 R₃ is a carbonyl containing moiety; and

R₂ is a linear or branched alkyl, alkylthio, alkylthioalkyl, cycloalkyl, alkylcycloalkyl, phenyl or a phenylalkyl optionally substituted with halogen, cyano, nitro, hydroxy, haloalkyl, alkylthio, terminal guanidine, guanidine, alkyl guanidine, dialkyl guanidine or amidine.

15

3. The compound according to claim 2 wherein R₃ is

20




4. The compound according to claim 3, wherein R₂ is 2-propyl, benzyl, 3-guanidinyl or 4-amidinylbenzyl.

25

5. The compound according to claim 4, of the formula


30



wherein

R_1 is an optionally substituted phenylalkyl; and
 R_2 is 2-propyl.




6. The compound according to claim 4, wherein R_1 is trifluoromethyl,
 5 methyl, difluoromethyl, 2,6-difluorobenzyl, benzyl, 3-thienylmethyl, styryl, 4-trifluoromethylstyryl, 4-methoxystyryl, 4-methoxybenzyl, phenyl, 2-phenylethyl, 3-methoxybenzyl, 3-phenylpropyl, 3-trifluoromethylbenzyl, 2-methoxybenzyl, or 2-trifluoromethylphenylchloromethyl.
- 10 7. The compound according to claim 5, wherein R_1 is 3-trifluoromethylbenzyl or 3-methoxybenzyl.
8. The compound according to claim 2 wherein R_3 is a moiety which contributes to the oral activity of said compound and is selected from the
 15 following:





(II)

where X_1 is N or CH;



- 20 X_2 is selected from the following:



25





30



where W is H or alkyl straight chained or branched;


X_3 is Ala, bAla, Leu, Ile, Val, Nva, bVal, Met, or Nle or an N-methyl derivative or a bond; and

- X_4 is an amino acid consisting of proline, isoleucine, leucine,
 5 cyclohexylalanine, cysteine optionally substituted at the sulfur with alkyl, alkenyl, being linear or branched or phenyl optionally substituted with halogen, cyano, nitro, haloalkyl, amino, aminoalkyl, dialkylamino, alkyl, alkoxy, haloalkoxy, carboxyl, carboalkoxy, alkylcarboxamide, arylcarboxamide, alkylthio, haloalkylthio groups being linear or branched, a
 10 phenylalanine, indoline-2-carboxylic acid tetrahydroisoquinoline-2-carboxylic acid optionally substituted with alkyl, alkenyl, haloalkenyl, alkynyl being linear or branched, halogen, cyano, nitro, haloalkyl, amino, aminoalkyl, dialkylamino, alkoxy, haloalkoxy, carbonyl, carbalkoxy, alkylcarboxamide, arylcarboxamide, alkylthio, or haloalkylthio group being linear or branched,
 15 tryptophan, valine, norvaline, norleucine, octahydroindole-2-carboxylic acid, lysine optionally substituted with alkylcarboxy or arylcarboxy, glycine optionally substituted at the nitrogen with alkyl, cycloalkyl, phenyl, bicycloalkyl, heteroaryl, alkenyl, alkynyl, cycloalkyl alkyl, bicycloalkyl alkyl, alkoxyalkyl, alkylthioalkyl, alkylaminoalkyl, fused aryl-cycloalkyl alkyl, fused
 20 heteroaryl-cycloalkyl, dialkylaminoalkyl, carboxyalkyl or alkoxy-carbonyl alkyl;
 or R_3 is



or

5



(v)

wherein Z_2 and Z_3 are optionally substituted alkyl, aryl, or arylalkyl;

10 Z_3 is optionally substituted alkyl, cycloalkyl, alkenyl, aryl, arylalkyl, aliphatic heterocycle, or aromatic heterocycle;

Z_4 and Z_6 are hydrogen or methyl; and

A is selected from the group consisting of


15



and n is 0, 1 or 2;

20 or R_3 is

25



where G is nitrogen or carbon optionally substituted with hydrogen, alkyl, cycloalkyl, aryl, heteroaryl or alkylheteroaryl;

J_1 and J_2 are independently selected from hydrogen, alkyl, cycloalkyl, 30 aryl, heteroaryl, or alkylaryl; and

J₃ is selected from hydrogen, alkylacyl, arylacyl, arylalkylacyl, alkyl-SO₂-, aryl-SO₂-, alkylaryl-SO₂-, heterocycle-SO₂-, alkyl-NH-SO₂-, aryl-NH-SO₂-, alkylaryl-NH-SO₂-, or heterocycle-NH-SO₂-.

- 5 9. A method of inhibiting neutrophil elastase which comprises administering to a host in need of such inhibition an effective amount of a compound according to claim 2.
- 10 10. A method of treating acute respiratory disease which comprises administering to a host in need of such treatment an effective amount of a compound according to claim 2.
- 15 11. A method of treating myocardial ischemia-reperfusion injury which comprises administering to a host in need of such treatment an effective amount of a compound according to claim 2.
12. A pharmaceutical composition comprising a compound according to claim 1 and a pharmaceutically acceptable carrier.




FIG. 1




FIG. 2




FIG. 3




FIG. 4

INTERNATIONAL SEARCH REPORT

International Application No

PCT/US 95/14989

A. CLASSIFICATION OF SUBJECT MATTER
IPC 6 C07K5/062 A61K38/05

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC 6 C07K A61K C07D

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	EP,A,0 291 234 (ICI AMERICA INC) 17 November 1988 see page 9, line 6 - page 10, line 34; claims; examples 12,14 ---	1-12
P,A	WO,A,95 21855 (ZENECA LTD ; PEGG STEPHEN JOHN (GB); SEPENDA GEORGE JOSEPH (GB); DA) 17 August 1995 see claims; examples ---	1-12
A	EP,A,0 480 044 (JAPAN TOBACCO INC) 15 April 1992 see page 2, line 8 - line 38; claims; examples 6,9 ---	1-12
A	EP,A,0 529 568 (MERRELL DOW PHARMA) 3 March 1993 see claims; examples 1-3; tables 2,3 --- -/--	8

☒ Further documents are listed in the continuation of box C.

☒ Patent family members are listed in annex.

* Special categories of cited documents:

- *A* document defining the general state of the art which is not considered to be of particular relevance
- *E* earlier document but published on or after the international filing date
- *L* document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
- *O* document referring to an oral disclosure, use, exhibition or other means
- *P* document published prior to the international filing date but later than the priority date claimed

- *T* later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
- *X* document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
- *Y* document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art
- *Z* document member of the same patent family

Date of the actual completion of the international search

29 March 1996

Date of mailing of the international search report

07. 05. 96

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2
NL - 2220 HV Rijswijk
Tel. (+ 31-70) 340-2040, Tx. 31 651 epo nl,
Fax: (+ 31-70) 340-3016

Authorized officer

Fuhr, C

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	GB,A,1 397 073 (GLAXO LAB LTD) 11 June 1975 see claims; examples ---	1
X	DE,A,22 24 338 (GLAXO LABORATORIES LTD.) 30 November 1972 see claims; examples -----	1

INTERNATIONAL SEARCH REPORT

International application No.

PCT/US95/14989

Box I Observations where certain claims were found unsearchable (Continuation of item 1 of first sheet)

This international search report has not been established in respect of certain claims under Article 17(2)(a) for the following reasons:

1. ☒ Claims Nos.: 9-11

because they relate to subject matter not required to be searched by this Authority, namely:

Remark: As far as these claims are directed to methods of treatment of the human/animal body, the search has been carried out and based on the alleged effects of the compound/composition.

- In view of -

2. ☐ Claims Nos.:
because they relate to parts of the international application that do not comply with the prescribed requirements to such an extent that no meaningful international search can be carried out, specifically:

3. ☐ Claims Nos.:
because they are dependent claims and are not drafted in accordance with the second and third sentences of Rule 6.4(a).

Box II Observations where unity of invention is lacking (Continuation of item 2 of first sheet)

This International Searching Authority found multiple inventions in this international application, as follows:

1. ☐ As all required additional search fees were timely paid by the applicant, this international search report covers all searchable claims.
2. ☐ As all searchable claims could be searched without effort justifying an additional fee, this Authority did not invite payment of any additional fee.
3. ☐ As only some of the required additional search fees were timely paid by the applicant, this international search report covers only those claims for which fees were paid, specifically claims Nos.:
4. ☐ No required additional search fees were timely paid by the applicant. Consequently, this international search report is restricted to the invention first mentioned in the claims; it is covered by claims Nos.:

Remark on Protest

☐ The additional search fees were accompanied by the applicant's protest.

☐ No protest accompanied the payment of additional search fees.

INTERNATIONAL SEARCH REPORT

International Application N. PCT/US95/14989

FURTHER INFORMATION CONTINUED FROM PCT/ISA/210

In view of the extremely large number of compounds falling under claims 1 and 12, and of the absence of any sensible support for these claims in the description, the International Searching Authority considers that it is not economically reasonable to draw up a report covering the entire subject matter of claims 1 and 12.

The search for claim 1 has therefore been limited and includes all the real examples given in the description and covers all similar compounds having the alleged activities as well (see also "remark").

INTERNATIONAL SEARCH REPORT

Inclusion on patent family members

International Application No

PCT/US 95/14989

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
EP-A-0291234	17-11-88	AU-B- 605284	10-01-91
		AU-B- 1539088	17-11-88
		CA-A- 1326107	11-01-94
		DE-D- 3888568	28-04-94
		DE-T- 3888568	21-07-94
		ES-T- 2063034	01-01-95
		FI-C- 91072	10-05-94
		IE-B- 62524	08-02-95
		IL-A- 86236	29-12-94
		JP-A- 1045395	17-02-89
		NO-B- 176519	09-01-95
		SU-A- 1678207	15-09-91
		US-A- 5164371	17-11-92
		ZA-A- 8803006	11-11-88
WO-A-9521855	17-08-95	AU-B- 1583995	29-08-95
EP-A-0480044	15-04-92	CA-A- 2056453	01-10-91
		WO-A- 9115487	17-10-91
		US-A- 5204462	20-04-93
EP-A-0529568	03-03-93	AU-B- 655831	12-01-95
		AU-B- 2106592	25-02-93
		CA-A- 2076307	23-02-93
		HU-A,B 62014	29-03-93
		JP-A- 5213991	24-08-93
		NZ-A- 243998	25-11-94
		US-A- 5478811	26-12-95
		ZA-A- 9206185	01-03-93
GB-A-1397073	11-06-75	BE-A- 783653	20-11-72
		DE-A- 2224338	30-11-72
		FR-A,B 2138079	29-12-72
		NL-A- 7206759	21-11-72
		US-A- 3991067	09-11-76
DE-A-2224338	30-11-72	GB-A- 1397073	11-06-75
		BE-A- 783653	20-11-72
		FR-A,B 2138079	29-12-72
		NL-A- 7206759	21-11-72

Patent document
cited in search reportPublication
datePatent family
member(s)Publication
date

DE-A-2224338

US-A- 3991067

09-11-76